arXiv:1303.2968v1 [math.PR] 12 Mar 2013

1D Log Gases and the Renormalized Energy : Crystallization at Vanishing Temperature

Etienne Sandier and Sylvia Serfaty

March 13, 2013

Abstract

We study the statistical mechanics of a one-dimensional log gas with general potential and arbitrary β, the inverse of temperature, according to the method we introduced for two-dimensional Coulomb gases in [SS2]. Such ensembles correspond to random matrix models in some particular cases. The formal limit $\beta = \infty$ corresponds to "weighted Fekete sets" and is also treated.

We introduce a one-dimensional version of the "renormalized energy" of [SS1], measuring the total logarithmic interaction of an infinite set of points on the real line in a uniform neutralizing background. We show that this energy is minimized when the points are on a lattice.

By a suitable splitting of the Hamiltonian we connect the full statistical mechanics problem to this renormalized energy W, and this allows us to obtain new results on the distribution of the points at the microscopic scale: in particular we show that configurations whose W is above a certain threshhold (which tends to $\min W$ as $\beta \to \infty$) have exponentially small probability. This shows that the configurations have increasing order and crystallize as the temperature goes to zero.

1 Introduction

In [SS2] we studied the statistical mechanics of a 2D classical Coulomb gas (or two-dimensional plasma) via the tool of the "renormalized energy" W introduced in [SS1], a particular case of which is the Ginibre ensemble in random matrix theory.

In this paper we are interested in doing the analogue in one dimension, i.e. first defining a "renormalized energy" for points on the real line, and applying this tool to the study of the classical log gases i.e. to probability laws of the form

$$d\mathbb{P}_n^\beta(x_1,\dots,x_n) = \frac{1}{Z_n^\beta} e^{-\frac{\beta}{2} w_n(x_1,\dots,x_n)} dx_1 \dots dx_n \tag{1.1}$$

where Z_n^β is the associated partition function, i.e. a normalizing factor such that $\mathbb{P}_n^\beta$ is a probability, and

$$w_n(x_1,\dots,x_n) = -\sum_{i\neq j} \log|x_i - x_j| + n\sum_{i=1}^n V(x_i). \tag{1.2}$$

Here the x_i's belong to $\mathbb{R}$, $\beta > 0$ is a parameter corresponding to (the inverse of) a temperature, and V is a relatively arbitrary potential, satisfying some growth conditions. For a general presentation, we refer to the textbook [For]. Minimizers of w_n are also called “weighted Fekete sets” and arise in interpolation, cf. [SaTo].

There is an abundant literature on the random matrix aspects of this problem (the connection was first pointed out by Wigner and Dyson [Wi, Dy]), which is the main motivation for studying log gases. Indeed, for the quadratic potential $V(x) = x^2/2$, particular cases of β correspond to the most famous random matrix ensembles: for $\beta = 1$ the law $\mathbb{P}_n^\beta$ is the law of eigenvalues of matrices of the Gaussian Orthogonal Ensemble (GOE), while for $\beta = 2$ it corresponds to the Gaussian Unitary Ensemble (GUE), for general reference see [For, AGZ, Me]. For $V(x)$ still quadratic, general β's have been shown to correspond to tri-diagonal random matrix ensembles, cf. [DE, ABF]. This observation allowed Valkó and Virág [VV] to derive the sine-β processes as the local spacing distributions of these ensembles. When $\beta = 2$ and $V(x)$ is more general, the model corresponds (up to minor modification) to other determinantal processes called orthogonal polynomial ensembles (see e.g. [Ko] for a review).

The study of $\mathbb{P}_n^\beta$ via the random matrix aspect is generally based on explicit formulas for correlation functions and local statistics, obtained via orthogonal polynomials, as pioneered by Gaudin, Mehta, Dyson, cf. [Me, D, DG]. We emphasize that for general β and V, which is the setting of the present work, there is no random matrix model associated and no explicit formulas available, and fewer results. Notable exceptions are the work of Ben Arous and Guionnet [BG] also valid for every β and the recent work of Bourgade, Erdös and Yau [BEY1, BEY2], which are in the same setting as ours. We will discuss them more below. The global and local statistics (e.g. spacings) of eigenvalues of large random matrices are expected to be independent of the fact that the entries are specifically Gaussian, this is refered to as “universality”, cf. [TV, EPRSY] for latest results. Here we are rather interested in questions of universality with respect to the potential V, as in [BEY1, BEY2].

The results here are counterparts of the results we obtained in [SS2] for $x_1, \dots, x_n$ belonging to $\mathbb{R}^2$ (this corresponds for V quadratic and $\beta = 2$ to the Ginibre ensemble of non-hermitian Gaussian random matrices), or in other words the two-dimensional Coulomb gas. The study in [SS2] relied on relating the Hamiltonian w_n to a Coulomb “renormalized energy” W introduced in [SS1] in the context of Ginzburg-Landau vortices. This relied crucially on the fact that the logarithm is the Coulomb kernel in two dimensions, or in other words the fundamental solution to the Laplacian. When looking at the situation in one dimension, i.e. the present situation of the 1D log-gas, the logarithmic kernel is no longer the Coulomb kernel, and it is not a priori clear that anything similar to the study in two dimensions can work. Note that the 1D Coulomb gas, corresponding to $\mathbb{P}_n^\beta$ where the logarithmic interaction is replaced by the 1D Coulomb kernel $|x|$, has been studied, notably by Lenard [Le1, Le2], Brascamp-Lieb [BL], Aizenman-Martin [AM]. The situation there is rendered again more accessible by the Coulomb nature of the interaction and its less singular character. In particular [BL] prove cristallization (i.e. that the points tend to arrange along a regular lattice) in the limit of a small temperature, we will get a similar result for the log-gas.

The starting point of our study is that even though the logarithmic kernel is not Coulombic in dimension 1, we can view the particles on the real line as embedded into the two-dimensional plane and interacting as Coulomb charges there. This provides a way of defining an analogue of the “renormalized energy” of [SS1] in the one-dimensional setting, still called W, which goes “via” the two-dimensional plane, see Section 1.1 below.

Once this is accomplished, we connect in the same manner as [SS2] the Hamiltonian w_n to the renormalized energy W via a "splitting formula" (cf. Lemma 1.7 below), and we obtain the counterparts results to [SS2]:

- a next-order expansion of the partition function in terms of n and β, cf. Theorem 3. For V quadratic, exact formulas were already known via Selberg integrals.
- the proof that ground states of w_n, or "weighted Fekete sets", converge to minimizers of W, cf. Theorem 4.
- the proof that the minimum of W is achieved by the one-dimensional regular lattice $\mathbb{Z}$, called the "clock distribution" in the context of orthogonal polynomial ensembles [Si]. This is in contrast with the dimension 2 where the identification of minimizers of W is still open (but conjectured to be "Abrikosov" triangular lattices.)
- A large deviations type result which shows that events with high W become less and less likely as $\beta \to \infty$, proving in particular the *crystallization* as the temperature tends to 0, in view of the result of the previous item.

As far as we know, this is the first proof of crystallization for one-dimensional log gases. Our renormalized energy W, which serves to prove the crystallization also appears (like its two-dimensional version) to be a measurement of "order" of a configuration at the microscopic scale $1/n$. What we show here is that there is more and more order (or rigidity) in the log gas, as the temperature gets small. Of course, as already mentioned, it is known that eigenvalues of random matrices, even of general Wigner matrices, should be regularly spaced. In fact the local statistics of the random process are completely known from [VV], and [BEY1, BEY2] showed that this could be extended to general V's. Our results approach this question sort of orthogonally, by exhibiting a unique number which measures the average rigidity, without using explicit formulas for local statistics relying on determinantal forms. Note that in [BSe] the second author and Borodin used W as a way of quantifying the order of random point processes, in particular those arising as local limits in random matrix theory.

Our study here differs technically from the two-dimensional one in two ways: the first one is in the definition of W by embedding the problem into the plane, as already mentioned. The second one is more subtle: in both settings a crucial ingredient in the analysis is to reduce the evaluation of the interactions to an extensive quantity (instead of sums of pairwise Coulomb interactions); that quantity is essentially the L^2 norm of the "electric field" generated by the Coulomb charges. Test-configurations can be built and their energy evaluated by "copying and pasting", provided a cut-off procedure is devised: it consists essentially in taking a given electric field and making it vanish on the boundary of a given box while not changing its energy too much. In physical terms, this corresponds to *screening* the field. The point is that screening is much easier in two dimensions than in one dimension, because in two dimensions there is more geometric flexibility to move charges around. The screening result of [SS2] does not apply to the 1-dimensional setting. We found that in fact, in dimension 1, not all configurations with finite energy can be effectively screened. However, we also found that generic "good" configurations can be, and this suffices for our purposes. The screening construction, which is different from the two-dimensional one, is one of the main difficulties here, and forms a large part of the paper.

The rest of the introduction is organized as follows: in Section 1.1, we present the embedding of the problem into two-dimensions and the definition of the one-dimensional renormalized energy W. In Section 1.2 we introduce notation and well-known facts on the equilibrium measure (i.e. the minimizer of the mean-field limiting Hamiltonian), as well as the "splitting formula" that connects w_n to W. Finally, in Section 1.3 we state our main results.

1.1 Definition of the renormalized energy in 1 dimension

As mentioned above, the renormalized energy between points in 1D is obtained by "embedding" the real line in the plane and computing the renormalized energy in the plane, as defined in [SS1]. More specifically, we introduce the following definitions:

$\mathbb{R}$ will denote the set of real numbers, but also the real line of the plane $\mathbb{R}^2$ i.e. points of the form $(x,0) \in \mathbb{R}^2$. For the sake of clarity, we will denote points in $\mathbb{R}$ by the letter x and points in the plane by $z=(x,y)$. For a function χ on $\mathbb{R}$, we define its natural extension $\bar{\chi}$ to a function on $\mathbb{R}^2$ by $\bar{\chi}(x,y) := \chi(x)$. I_R will denote the intervals $[-R/2, R/2]$ in $\mathbb{R}$ and $\bar{I}_R$ the strips $I_R \times \mathbb{R}$ in $\mathbb{R}^2$. $\delta_{\mathbb{R}}$ denotes the measure of length on $\mathbb{R}$ seen as embedded in $\mathbb{R}^2$, that is

$$\int_{\mathbb{R}^2} \varphi \delta_{\mathbb{R}} = \int_{\mathbb{R}} \varphi(x,0)\, dx$$

for any smooth compactly supported test function φ in $\mathbb{R}^2$. This measure can be multiplied by bounded functions on the real-line. For any measurable set U, $|U|$ denotes its Lebesgue measure, and if U is a finite set $\#U$ denotes its cardinal. δ_p denotes the Dirac mass at p.

The renormalized energy W is an energy of an infinite configuration of points on the real line, screened by a uniform background charge, again on the real line. It will later be obtained as a limit as $n \to \infty$ of logarithmic interactions of systems of n particles/points.

We first introduce the classes $\mathcal{A}_m$ corresponding to infinite configurations on the real line with density m.

Definition 1.1. *Let m be a nonnegative number. Let E be a vector field in $\mathbb{R}^2$. We say E belongs to the admissible class $\mathcal{A}_m$ if*

$$\text{div } E = 2\pi(\nu - m\delta_{\mathbb{R}}), \quad \text{curl}\, E = 0 \quad \text{in } \mathbb{R}^2 \tag{1.3}$$

where ν has the form

$$\nu = \sum_{p\in\Lambda} \delta_p \quad \text{for some discrete set } \Lambda \subset \mathbb{R} \subset \mathbb{R}^2, \tag{1.4}$$

and

$$\frac{\nu(I_R)}{R} \quad \text{is bounded by a constant independent of } R>1. \tag{1.5}$$

Definition 1.2. *Let m be a nonnegative number. For any bounded function χ and any $E \in \mathcal{A}_m$ satisfying* (1.3) *we let*

$$W(E,\chi) = \lim_{\eta\to 0} \left(\frac{1}{2} \int_{\mathbb{R}^2 \setminus \cup_{p\in\Lambda} B(p,\eta)} \bar{\chi}|E|^2 + \pi \log \eta \sum_{p\in\Lambda} \chi(p) \right) \tag{1.6}$$

where $\bar{\chi}$ is the natural extension of χ.

Here we thus look at any configuration of points in the plane through an "electric field" E that it generates *in the whole plane* $\mathbb{R}^2$. Observe that if E satisfies (1.3), there exists a function H such that $E = -\nabla H$, we will call this function the *potential* generated by the point charges. We also note that the distribution of charges on the real line is compensated by a background charge $m\delta_{\mathbb{R}}$ which is also concentrated on the real line. We will use the notation χ_R for positive cutoff functions over $\mathbb{R}$ satisfying, for some constant C independent of R,

$$|\nabla \chi_R| \le C, \quad \mathrm{Supp}(\chi_R) \subset I_R, \quad \chi_R(x) = 1 \text{ if } |x| < R/2 - 1. \tag{1.7}$$

Definition 1.3. *The renormalized energy W is defined, for $E \in \mathcal{A}_m$, by*

$$W(E) = \limsup_{R\to\infty} \frac{W(E,\chi_R)}{R}, \tag{1.8}$$

where $\{\chi_R\}_{R>0}$ satisfies (1.7).

Returning to writing $E = -\nabla H$, we see that while W in 2D can be viewed as a "renormalized" way of computing $\|H\|_{H^1(\mathbb{R}^2)}$, in 1D it amounts rather to a renormalized computation of $\|H\|_{H^{1/2}(\mathbb{R})}$ (where H^s denote the fractional Sobolev spaces). In other words, because the logarithmic kernel is not Coulombic in one-dimension, the associated energy is non-local (and the associated operator is the fractional Laplacian $\Delta^{1/2}$). Augmenting the dimension by 1 allows to make it local and Coulombic again. This well-known extension idea is also now commonly used in the study of general fractional Laplacians [CS].

As in the two dimensional case, we have the following properties:

- The value of W does not depend on $\{\chi_R\}_R$ as long as it satisfies (1.7).
- W is insensitive to compact perturbations of the configuration.
- Scaling: it is easy to check that if E belongs to $\mathcal{A}_m$ then $E' := \frac{1}{m}E(\cdot/m)$ belongs to $\mathcal{A}_1$ and

  $$W(E) = m\left(W(E') - \pi \log m\right). \tag{1.9}$$

- If $E \in \mathcal{A}_m$ then in the neighborhood of $p \in \Lambda$ we have $\operatorname{div} E = 2\pi(\delta_p - m\delta_{\mathbb{R}})$, $\operatorname{curl} E = 0$, thus we have near p the decomposition $E(x) = -\nabla \log|x-p| + f(x)$ where f is smooth, and it easily follows that the limit (1.6) exists. It also follows that E belongs to L^q_{loc} for any $q < 2$.

An extra fact proven below is valid only in the one-dimensional case (and not the two-dimensional one):

Lemma 1.4. *Let $E \in \mathcal{A}_m$ be such that $W(E) < +\infty$. Then any other E' satisfying* (1.3)–(1.4) *with the same ν and $W(E') < +\infty$, is such that $E' = E$. In other words, W only depends on the points.*

By simple considerations similar to [SS2, Section 1.2] this makes W a measurable function of the bounded Radon measure ν.

This implies in particular

Corollary 1.5. *Under the same assumptions, if* $S(x,y)=(x,-y)$ *then* $E\circ S=S\circ E$.

Indeed, it is easy to check that $E'=S\circ E\circ S$ satisfies (1.3) with the same ν as E, and obviously $W(E')<+\infty$, hence $E'=E$.

The following lemma is proven in [BSe], and shows that there is an explicit formula for W in terms of the points when the configuration is assumed to have some periodicity. Here we can reduce to $m=1$ by scaling, as seen above.

Lemma 1.6. *In the case* $m=1$ *and when the set of points* Λ *is periodic with respect to some lattice* $N\mathbb{Z}$*, then it can be viewed as a set of* N *points* $a_1,\dots,a_N$ *over the torus* $\mathbb{T}_N:=\mathbb{R}/(N\mathbb{Z})$*. In this case, by Lemma 1.4 there exists a unique* E *satisfying* (1.3) *and for which* $W(E)<+\infty$*. It is periodic and equal to* $E_{\{a_i\}}=\nabla H$*, where* H *is the solution on* $\mathbb{T}_N$ *to* $-\Delta H=2\pi(\sum_i \delta_{a_i}-\delta_{\mathbb{R}})$*, and we have the explicit formula:*

$$W(E_{\{a_i\}})=-\frac{\pi}{N}\sum_{i\neq j}\log\left|2\sin\frac{\pi(a_i-a_j)}{N}\right|-\pi\log\frac{2\pi}{N}. \tag{1.10}$$

As in the two-dimensional case, we can prove that $\min_{\mathcal{A}_m}$ is achieved, but contrarily to the two-dimensional case, the value of the minimum can be explicitly computed: we will prove here the following

Theorem 1. $\min_{\mathcal{A}_m} W=-\pi m\log(2\pi m)$ *and this minimum is achieved by the perfect lattice i.e.* $\Lambda=\frac{1}{m}\mathbb{Z}$*.*

We recall that in dimension 2, it was conjectured in [SS1] but not proven, that the minimum value is achieved at the triangular lattice with angles 60° (which is shown to achieve the minimum among all lattices), also called the Abrikosov lattice in the context of superconductivity.

The proof of Theorem 1 relies on showing that a minimizer can be approximated by configurations which are periodic with period $N\to\infty$ (this result itself relies on the screening construction we mentioned above), and then using a convexity argument to find the minimizer among periodic configurations with a fixed period via (1.10).

1.2 The equilibrium measure and the splitting formula

The Hamiltonian (1.2) is written in the mean-field scaling. The limiting "mean-field" limiting energy and its minimizer are in fact well-known, and they tell us the optimal *macroscopic* distribution of points — what we investigate here is the *microscopic* distribution of points, as a correction to this optimal macroscopic behavior. More precisely, the mean-field limit energy is

$$\mathcal{F}(\mu)=\int_{\mathbb{R}\times\mathbb{R}}-\log|x-y|\,d\mu(x)\,d\mu(y)+\int_{\mathbb{R}}V(x)\,d\mu(x), \tag{1.11}$$

and its unique minimizer among probability measures is called the *equilibrium measure* in the language of potential theory (cf. [SaTo]). We will denote it μ_0. It is not hard to prove that the "spectral measure" (so-called in the context of random matrices) $\nu_n=\frac{1}{n}\sum_{i=1}^n\delta_{x_i}$ converge to μ_0, the minimizer of $\mathcal{F}$, itself also called Voiculescu's noncommutative entropy in the context of random matrices, cf. e.g. [AGZ] and references therein. The sense of convergence usually proven is

$$\mathbb{P}_n^\beta\left(\forall f\in C_b(\mathbb{C},\mathbb{R}), \int f\,d\nu_n \to \int f\,d\mu_0\right)=1 \tag{1.12}$$

For example, for the case of the GUE i.e. when $V(x)=|x|^2$ and $\beta=1$, the correspond distribution μ_0 is simply Wigner's "semi-circle law" $\rho(x)=\frac{1}{2\pi}\sqrt{4-x^2}\mathbf{1}_{|x|<2}$, cf. [Wi, Me]. A stronger result was proven in [BG] for all β (cf. [AGZ] for the case of general V): it estimates the large deviations from this convergence and shows that $\mathcal{F}$ is the appropriate rate function. The result can be written:

Theorem 2 (Ben Arous - Guionnet [BG])**.** *Let $\beta>0$, and denote by $\widetilde{\mathbb{P}}_n^\beta$ the image of the law (1.1) by the map $(x_1,\dots,x_n)\mapsto \nu_n$, where $\nu_n=\frac{1}{n}\sum_{i=1}^n\delta_{x_i}$. Then for any subset A of the set of probability measures on $\mathbb{R}$ (endowed with the topology of weak convergence), we have*

$$-\inf_{\mu\in\mathring{A}}\widetilde{\mathcal{F}}(\mu)\le \liminf_{n\to\infty}\frac{1}{n^2}\log\widetilde{\mathbb{P}}_n^\beta(A)\le \limsup_{n\to\infty}\frac{1}{n^2}\log\widetilde{\mathbb{P}}_n^\beta(A)\le -\inf_{\mu\in\bar{A}}\widetilde{\mathcal{F}}(\mu),$$

where $\widetilde{\mathcal{F}}=\mathcal{F}-\min\mathcal{F}$.

Our result Theorem 5 below will be a sort of next-order extension of this result. Note that the Central Limit Theorem for (macroscopic) fluctuations from the limiting law was proved by Johansson [Jo].

Let us now state a few facts that we will need about the equilibrium measure μ_0, for which we refer to [SaTo]: μ_0 is characterized by the fact that there exists a constant c (depending on V) such that

$$U^{\mu_0}+\frac{V}{2}=c \text{ quasi-everywhere in the support of } \mu_0, \text{ and } U^{\mu_0}+\frac{V}{2}\ge c \text{ quasi-everywhere} \tag{1.13}$$

where for any μ, U^μ is the potential generated by μ, defined by

$$U^\mu(x)=-\int_{\mathbb{R}}\log|x-y|\,d\mu(y). \tag{1.14}$$

(Note that looking in Fourier, this corresponds to $U^\mu=-\Delta^{-1/2}\mu$). Here and in all the paper, we denote by Δ^{-1} the operator of convolution by $\frac{1}{2\pi}\log|\cdot|$ in $\mathbb{R}^2$. It is such that $\Delta\circ\Delta^{-1}=Id$, where Δ is the usual Laplacian. We may view μ_0 as a measure over $\mathbb{R}^2$ and extend U^μ into a function of $\mathbb{R}^2$ by $U^\mu(z)=-\int_{\mathbb{R}}\log|z-(y,0)|\,d\mu(y)$. Then $U^\mu=-\Delta^{-1}\mu$ in $\mathbb{R}^2$.

We also define

$$\zeta=U^{\mu_0}+\frac{V}{2}-c$$

where c is the constant in (1.13). From the above we know that $\zeta\ge0$ in $\mathbb{R}$ and $\zeta=0$ in $\Sigma:=Supp(\mu_0)$. We will make the assumption that μ_0 has a density m_0 with respect to the Lebesgue measure.

It's now time to state our assumptions on V: we assume in the sequel

$$V \text{ is lower semicontinuous and } \lim_{|x|\to+\infty}\frac{V(x)}{2}-\log|x|=+\infty. \tag{1.15}$$

$$\Sigma \text{ is a finite union of closed intervals } \Sigma_1, \dots, \Sigma_M. \tag{1.16}$$

$$\text{There exist } \gamma, \overline{m} > 0 \text{ such that } \gamma\sqrt{\text{dist } (x, \mathbb{R} \setminus \Sigma)} \le m_0(x) \le \overline{m} \text{ for all } x \in \mathbb{R}. \tag{1.17}$$

$$m_0 \in C^{0,\frac{1}{2}}(\mathbb{R}). \tag{1.18}$$

$$\text{There exists } \beta_1 > 0 \text{ such that } \int_{\mathbb{R}\setminus[-1,1]} e^{-\beta_1(V/2(x) - \log|x|)}\, dx < +\infty. \tag{1.19}$$

The assumption (1.15) ensures (see [SaTo]) that (1.11) has a minimizer, and that its support Σ is compact. Assumptions (1.16)–(1.18) are needed for the construction in Section 4. They are certainly not optimal but are meant to include at least the model case of $\mu_0 = \rho$, Wigner's semi-circle law. Assumption (1.19) is a supplementary assumption on the growth of V at infinity, needed for the case with temperature. It only requires a very mild growth of $V/2 - \log|x|$, i.e. slightly more than (1.15).

We are interested in examining the "next order" behavior, or that of fluctuations around the limiting law μ_0. While $\mathcal{F}$ and μ_0 are found through the large deviations at speed n^2, as seen in Theorem 2, we look into the speed n and show, not a large deviations principle, but that a suitably defined average $\int W\, dP$ of the "renormalized energy" W defined above (1.8) acts as a sort of rate function, and that a "threshhold phenomenon" holds: $\int W\, dP \le \min W + C_\beta$ except with exponentially small probability. The crucial fact is that $C_\beta \to 0$ as $\beta \to \infty$, which shows the crystallization. The average of W corresponds to the average of W computed over all blown-up configurations of the points (with respect to all possible blow-up centers in Σ) at the scale n. The measure P is like a Young measure on all the possible limits of blow-ups of E.

The starting point of our analysis is to establish, through a simple but crucial exact "splitting formula", a link between the law $\mathbb{P}_n^\beta$ and the energy w_n, and W:

Lemma 1.7 (Splitting formula)**.** *For any $x_1, \dots, x_n \in \mathbb{R}$ the following holds*

$$w_n(x_1, \dots, x_n) = n^2\mathcal{F}(\mu_0) - n\log n + \frac{1}{\pi} W(\nabla H_n', \mathbf{1}_{\mathbb{R}^2}) + 2n\sum_{i=1}^n \zeta(x_i) \tag{1.20}$$

where H_n' is the solution in $\mathbb{R}^2$ to

$$H_n'(x') = -2\pi\Delta^{-1}\left(\sum_{i=1}^n \delta_{x_i'} - m_0(x'/n)\delta_{\mathbb{R}}\right) \tag{1.21}$$

with $x_i' = nx_i$ (viewed as points in $\mathbb{R}^2$), and $W(\cdot,\cdot)$ is as in (1.6)*.*

We may then define

$$F_n(\nu) = \begin{cases} \frac{1}{n}\left(\frac{1}{\pi}W(\nabla H_n', \mathbf{1}_{\mathbb{R}^2}) + 2n\int_{\mathbb{R}} \zeta\, d\nu\right) & \text{if } \nu \text{ is of the form } \sum_{i=1}^n \delta_{x_i} \\ +\infty & \text{otherwise} \end{cases} \tag{1.22}$$

and also

$$\widehat{F_n}(\nu) = F_n(\nu) - 2\int_{\mathbb{R}} \zeta\, d\nu \le F_n(\nu) \tag{1.23}$$

and we thus have the following rewriting of w_n:

$$\boxed{w_n(x_1,\dots,x_n) = n^2\mathcal{F}(\mu_0) - n\log n + nF_n(\nu).} \tag{1.24}$$

This allows to separate orders in the limit $n\to\infty$ since one of the main outputs of our analysis is that $F_n(\nu)$ is of order 1. Moreover, $F_n(\nu)$ contains information about the microscopic structure of the configurations, its limit as $n\to\infty$ will be the average of W over all blown-up points, $\int W\,dP$ mentioned above.

1.3 Main results

The first main result is the announced one on the next order asymptotic expansion of the partition function, which becomes sharp as $\beta\to\infty$.

Theorem 3. *Let V satisfy assumptions* (1.15)—(1.19)*. There exist functions f_1, f_2 depending only on V, such that for any $\beta_0>0$ and any $\beta\ge\beta_0$, and for n larger than some n_0 depending on β_0, we have*

$$n\beta f_1(\beta)\le \log Z_n^\beta - \left(-\frac{\beta}{2}n^2\mathcal{F}(\mu_0)+\frac{\beta}{2}n\log n\right)\le n\beta f_2(\beta), \tag{1.25}$$

with f_1, f_2 bounded in $[\beta_0,+\infty)$ and

$$\lim_{\beta\to\infty} f_1(\beta)=\lim_{\beta\to\infty} f_2(\beta)=\frac{\alpha}{2} \tag{1.26}$$

where

$$\alpha=\int_\Sigma \mu_0(x)\log(2\pi\mu_0(x))\,dx. \tag{1.27}$$

Remark 1.8. *In fact we prove that the statement holds with $f_2(\beta)=\frac{\alpha}{2}+\frac{C}{\beta}$ for any $C>\log|\Sigma|$.*

The exact value for Z_n^β is only known for arbitrary β's for $V(x)=x^2/2$ via Selberg integrals cf. [Me, AGZ]. For some more general potentials there are formulas for $\beta=2$ (e.g. [EML, PAS, GMS]). For general potential and β, this improves on the known results, which only gave the expansion $\log Z_n^\beta\sim\beta n^2\mathcal{F}(\mu_0)$.

Our next result is the one we mentioned about minimizers of w_n, or weighted Fekete sets. It identifies the Γ-limit (in the sense of Γ-convergence, cf. [Br]) of $\{F_n\}_n$, defined in (1.22) or (1.24). This allows a description of the minimizers at the microscopic level. Below we abuse notation by writing $\nu_n=\sum_{i=1}^n\delta_{x_i}$ when it should be $\nu_n=\sum_{i=1}^n\delta_{x_{i,n}}$. For such a ν_n we let $\nu_n'=\sum_{i=1}^n\delta_{x_i'}$ be the measure in blown-up coordinates $x'=nx$, and $E_{\nu_n}=-\nabla H_n'$, where H_n' is defined by (1.21). (To avoid confusion, we emphasize here that ν_n lives at the original scale while E_{ν_n} lives at the blown-up scale.) We set $m_0{}'(x')=m_0(x'/n)$, where m_0 is the density of μ_0. We also let

$$P_{\nu_n}=\fint_\Sigma \delta_{(x,E_{\nu_n}(nx+\cdot))}\,dx, \tag{1.28}$$

i.e. the push-forward of the normalized Lebesgue measure on Σ by $x\mapsto(x,E_{\nu_n}(nx+\cdot))$. It is a probability measure on $\Sigma\times L^p_{\text{loc}}(\mathbb{R}^2,\mathbb{R}^2)$ (couples of (blow-up centers, blown-up electric field around this center)). In taking the Γ-limit of F_n, the limiting object is more complex, it is the limit P of P_{ν_n}, i.e. a Young measure akin to the Young measures on micropatterns introduced in [AM]. Note that in the rest of the paper, the probability P has nothing to do with $\mathbb{P}_n^\beta$, and depends on the realizations of the configurations of points. We state the generalized invariance property satisfied by these limiting object as a definition.

Definition 1.9. *We say a probability measure P on $\Sigma \times L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ is $T_{\lambda(x)}$-invariant if P is invariant by $(x, E) \mapsto (x, E(\lambda(x) + \cdot))$, for any $\lambda(x)$ of class C^1 from Σ to $\mathbb{R}$.*

Theorem 4 (Microscopic behavior of weighted Fekete sets)**.** *Let the potential V satisfy assumptions* (1.15)*–*(1.18)*. Fix $1 < p < 2$ and let $X = \Sigma \times L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$.*

A. Lower bound. *Let $\nu_n = \sum_{i=1}^n \delta_{x_i}$ be a sequence such that $\widehat{F_n}(\nu_n) \le C$. Then P_{ν_n} defined by* (1.28) *is a probability measure on X and*

1. *Any subsequence of $\{P_{\nu_n}\}_n$ has a convergent subsequence converging to a probability measure on X as $n \to \infty$. We denote by P such a limit.*
2. *The first marginal of P is the normalized Lebesgue measure on Σ. P is $T_{\lambda(x)}$-invariant.*
3. *For P almost every (x, E) we have $E \in \mathcal{A}_{m_0(x)}$.*
4. *It holds that*

$$\liminf_{n\to\infty} \widehat{F_n}(\nu_n) \ge \frac{|\Sigma|}{\pi} \int W(E)\, dP(x, E) \ge \alpha, \tag{1.29}$$

where α is as in (1.27)*.*

B. Upper bound construction. *Conversely, assume P is a $T_{\lambda(x)}$-invariant probability measure on X whose first marginal is $\frac{1}{|\Sigma|} dx_{|\Sigma}$ and such that for P-almost every (x, E) we have $E \in \mathcal{A}_{m_0(x)}$. Then there exists a sequence $\{\nu_n = \sum_{i=1}^n \delta_{x_i}\}_n$ of empirical measures on Σ and a sequence $\{E_n\}_n$ in $L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that* div *$E_n = 2\pi(\nu_n' - m_0{}'\delta_{\mathbb{R}})$ and such that defining P_n as in* (1.28)*, with E_n replacing E_{ν_n}, we have $P_n \to P$ as $n \to \infty$ and*

$$\limsup_{n\to\infty} F_n(\nu_n) \le \frac{|\Sigma|}{\pi} \int W(E)\, dP(x, E). \tag{1.30}$$

C. Consequences for minimizers. *If $(x_1, \dots, x_n)$ minimizes w_n for every n and $\nu_n = \sum_{i=1}^n \delta_{x_i}$, then the limit P of P_{ν_n} as defined in* (1.28) *satisfies the following.*

1. *For P-almost every (x, E), the electric field E minimizes W over $\mathcal{A}_{m_0(x)}$.*
2. *We have*

$$\lim_{n\to\infty} F_n(\nu_n) = \lim_{n\to\infty} \widehat{F_n}(\nu_n) = \frac{|\Sigma|}{\pi} \int W(E)\, dP(x, E) = \alpha, \quad \lim_{n\to\infty} \sum_{i=1}^n \zeta(x_i) = 0.$$

Note that part B of the theorem is only a partial converse to part A because the constructed E_n need not be curl-free, hence in general $E_n \neq E_{\nu_n}$.

It can be expected that ζ (which is positive exactly in the complement of Σ) controls the distance to Σ (to some power). This can probably be proven under some strict growth assumption on V and explicit formulas for the Poisson kernel of the half-Laplacian. We do not pursue this question here.

The next result is the announced next order result on deviations, to be compared to Theorem 2.

We let A_n be a subset of $\mathbb{R}^n$. We identify points in $\mathbb{R}^n$ with measures ν_n of the form $\sum_{i=1}^n \delta_{x_i}$. We also embed $\mathbb{R}^n$ into the set of probabilities on $X = \Sigma \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ in the following way: For any n and $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ we let $i_n(\mathbf{x}) = P_{\nu_n}$, where $\nu_n = \sum_{i=1}^n \delta_{x_i}$ and P_{ν_n} is as in (1.28), so that $i_n(\mathbf{x})$ is an element of $\mathcal{P}(X)$, the set of probability measures on $X = \Sigma \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$, which we consider endowed with the topology of weak convergence.

Theorem 5. *Let V satisfy (1.15)–(1.19). For any $\beta_0 > 0$ and any $\beta \geq \beta_0$ the following holds.*

A. *For any $n > 0$, let $A_n \subset \mathbb{R}^n$. Denote*

$$A_\infty = \bigcap_{n>0} \overline{\bigcup_{m>n} i_m(A_m)}. \tag{1.31}$$

Then for any $\eta > 0$ there is $C_\eta > 0$ depending on V and η only such that α being as in (1.27),

$$\limsup_{n\to\infty} \frac{\log \mathbb{P}_n^\beta(A_n)}{n} \leq -\frac{\beta}{2}\Big(\frac{|\Sigma|}{\pi} \inf_{P \in A_\infty} \int W(E)\, dP(x,E) - \alpha - \eta - \frac{C_\eta}{\beta}\Big). \tag{1.32}$$

Conversely, let $A \subset \mathcal{P}(X)$ be a set of $T_{\lambda(x)}$-invariant probability measures on X and let $\mathring{A}$ be the interior of A for the topology of weak convergence. Then for any $\eta > 0$, there exists a sequence of subsets $A_n \subset \Sigma^n$ such that

$$-\frac{\beta}{2}\left(\frac{|\Sigma|}{\pi} \inf_{P \in \mathring{A}} \int W(E)\, dP(x,E) - \alpha + \eta + \frac{C_\eta}{\beta}\right) \leq \liminf_{n\to\infty} \frac{\log \mathbb{P}_n^\beta(A_n)}{n}, \tag{1.33}$$

and such that for any sequence $\{\nu_n = \sum_{i=1}^n \delta_{x_i}\}_n$ such that $(x_1, \dots, x_n) \in A_n$ for every n there exists a sequence of fields $E_n \in L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that $\operatorname{div} E_n = 2\pi(\nu_n' - m_0' \delta_{\mathbb{R}})$ *and such that — defining P_n as in (1.28) with E_n replacing E_{ν_n} — we have*

$$\lim_n P_n \in \mathring{A}. \tag{1.34}$$

B. *For each integer n, and given $\beta \geq \beta_0$ let $\mathbb{Q}_n^\beta$ denote the push-forward of $\mathbb{P}_n^\beta$ by i_n defined above. Then $\mathbb{Q}_n^\beta$ is tight in $\mathcal{P}(\mathcal{P}(X))$ and converges, up to a subsequence, to a probability measure $\mathbb{Q}_\beta \in \mathcal{P}(\mathcal{P}(X))$.*

Note that if P_n was P_{ν_n}, then (1.34) would be equivalent to saying that $\cap_n \overline{\cup_{m>n} i(A_m)} \subset \mathring{A}$. The difference between P_{ν_n} and P_n is that the latter is generated by a field E_n which is not necessarily curl free (we believe this gap cannot be bridged).

Part B of the result shows the existence of a limiting "electric field" process, hence equivalently, via projecting by $(x, E) \mapsto \frac{1}{2\pi} \operatorname{div} E + m_0(x)\delta_{\mathbb{R}}$, of a limiting point process.

As we said, Part A of this theorem means that events such that the average of W is larger than $\alpha + \eta + C_\eta/\beta$ have exponentially decaying probability as $n \to \infty$, with $\eta > 0$ arbitrary. Note here the important fact that

$$\min \frac{|\Sigma|}{\pi} \int W(E)\, dP(x,E) = \alpha \tag{1.35}$$

where the min is taken over P probabilities satisfying P-a.e. (x,E) satisfies $E \in \mathcal{A}_{m_0(x)}$, a fact which simply follows from (1.9) and Theorem 1.

This means that configurations should have more and more order (evaluated via W) as β gets large, and have to crystallize to minimizers of W when $\beta \to \infty$ (the term crystallization can be legitimately used in view of Theorem 1). Note that we do not claim that this is a true large deviations result, since the corresponding lower bound is missing. In fact it is likely that the large deviations rate function is not truly W but contains an extra entropy term. To demonstrate its existence is an open question.

Finally, let us mention that our method yields estimates on the probability of some rare events, typically the probability that the number of points in a microscopic interval deviates from the number given by μ_0. We present them below, even though stronger results are obtained in [BEY1, BEY2]. The results below follow easily from the estimate (provided by Theorem 3) that $\widehat{F_n} \le C$ except on a set of small probability.

Theorem 6. *Let V satisfy assumptions* (1.15)–(1.18). *There exists a universal constant $R_0 > 0$ and $c, C > 0$ depending only on V such that: For any $\beta_0 > 0$, any $\beta \ge \beta_0$, any n large enough depending on β_0, for any $x_1, \dots, x_n \in \mathbb{R}$, any $R > R_0$, any interval $I \subset \mathbb{R}$ of length R/n, and any $\eta > 0$, letting $\nu_n = \sum_{i=1}^n \delta_{x_i}$, we have the following:*

$$\log \mathbb{P}_n^\beta \left(|\nu_n(I) - n\mu_0(I)| \ge \eta R\right) \le -c\beta \min(\eta^2, \eta^3) R^2 + C\beta(R+n) + Cn, \tag{1.36}$$

$$\log \mathbb{P}_n^\beta \left((1 + R^2/n^2)^{\frac{1}{2} - \frac{1}{q}} \|\nu_n - n\mu_0\|_{W^{-1,q}(I)} \ge \eta\sqrt{n} \right) \le -cn\beta\eta^2 + Cn(\beta+1) \tag{1.37}$$

where $W^{-1,q}(I)$ is the dual of the Sobolev space $W_0^{1,q'}(I)$, with $1/q + 1/q' = 1$, in particular $W^{-1,1}$ is the dual of Lipschitz functions; and

$$\log \mathbb{P}_n^\beta \left(\int \zeta \, d\nu_n \ge \eta \right) \le -\frac{1}{2} n\beta\eta + Cn(\beta+1). \tag{1.38}$$

Note that in these results R can be taken to depend on n.

(1.36) tells us that the density of eigenvalues is correctly approximated by the limiting law μ_0 at all small scales bigger than $Cn^{-1/2}$ for some C. However this in fact should hold at all scales with $R \gg 1$, cf. [ESY, BEY1, BEY2]. (1.38) serves to control the probability that points are outside Σ (since $\{\zeta > 0\} = \Sigma^c$).

The rest of the paper is organized as follows. In Section 2 we prove the preliminary results Lemma 1.4 and 1.7, as well as Theorem 1. In Section 3 we turn to the core of the proof, with the statement of the main screening result and construction. Sections 4 and 5 contains the proofs. Finally Section 6 gathers the proofs of all the remaining results which are adapted from [SS2].

Acknowledgements: We are grateful to Alexei Borodin, Gérard Ben Arous, Amir Dembo, Percy Deift, Nicolas Fournier, Alice Guionnet and Ofer Zeitouni for their interest and helpful discussions. E. S. was supported by the Institut Universitaire de France and S.S. by a EURYI award.

2 The first results

2.1 Proof of Lemma 1.4

Assume E and E' belong to $\mathcal{A}_m$ and satisfy (1.3) with the same ν. Then $f = E - E'$ is divergence-free and curl-free, hence can be seen, identifying $\mathbb{R}^2$ and $\mathbb{C}$, as an entire holomorphic function $\sum_{n=0}^{\infty} a_n z^n$. If we assume that $W(E)$ and $W(E')$ are finite, then it follows from [SeTi], Corollary 1.2 that for $p < 2$, the growth of the L^1 norms of E and E' is no worse than $R^{3/2}\sqrt{\log R}$ hence there exists $C > 0$ such that for any $R > 2$ we have $\|f\|_{L^1(B_R)} \le CR^{3/2}\sqrt{\log R}$. But by Cauchy's formula we have, for any $R > 0$ and $t \in [R, R+1]$

$$a_n = \frac{1}{2i\pi} \int_{\partial B(0,t)} \frac{f(z)}{z^{n+1}}\, dz = \frac{1}{2i\pi} \int_R^{R+1} \int_{\partial B(0,t)} \frac{f(z)}{z^{n+1}}\, dz.$$

It follows with the above that $|a_n| \le CR^{3/2}\sqrt{\log R} R^{-n-1}$ which implies, letting $R \to \infty$ that $a_n = 0$ for any $n \ge 1$, thus f is a constant. This constant must then be zero since both E and E' are square integrable on the infinite strips $[a, b] \times [1, +\infty]$.

2.2 Proof of the splitting formula (Lemma 1.7)

Let $\nu_n = \sum_{i=1}^n \delta_{x_i}$. First, letting $\triangle$ denote the diagonal of $\mathbb{R} \times \mathbb{R}$, we may rewrite w_n as

$$w_n(x_1, \dots, x_n) = \int_{\triangle^c} -\log|x-y|\, d\nu_n(x)\, d\nu_n(y) + n \int_{\mathbb{R}} V(x)\, d\nu_n(x).$$

Splitting ν_n as $n\mu_0 + \nu_n - n\mu_0$ and using the fact that $\mu_0 \times \mu_0(\triangle) = 0$, we obtain

$$\begin{aligned} w(x_1, \dots, x_n) = n^2 \mathcal{F}(\mu_0) + 2n \int U^{\mu_0}(x)\, d(\nu_n - n\mu_0)(x) + n \int V(x)\, d(\nu_n - n\mu_0)(x) \\ + \int_{\triangle^c} -\log|x-y|\, d(\nu_n - n\mu_0)(x)\, d(\nu_n - n\mu_0)(y). \end{aligned}$$

Since $U^{\mu_0} + \frac{V}{2} = c + \zeta$ and since ν_n and $n\mu_0$ have same mass n, we have

$$2n \int U^{\mu_0}(x)\, d(\nu_n - n\mu_0)(x) + n \int V(x)\, d(\nu_n - n\mu_0)(x) = 2n \int \zeta\, d(\nu_n - \mu_0) = 2n \int \zeta\, d\nu_n,$$

using the fact that $\zeta = 0$ on the support of μ_0.

In addition, we have that

$$\int_{(\mathbb{R}\times\mathbb{R})\setminus\triangle} -\log|x-y|\, d(\nu_n - n\mu_0)(x)\, d(\nu_n - n\mu_0)(y) = \frac{1}{\pi} W(\nabla H_n, \mathbf{1}_{\mathbb{R}^2}), \tag{2.1}$$

where we define $H_n = -2\pi\Delta^{-1}\left(\sum_{i=1}^n \delta_{x_i} - n\mu_0\right)$. Indeed, the integral might as well be written as over $\mathbb{R}^2 \times \mathbb{R}^2 \setminus \triangle$ with the diagonal in $\mathbb{R}^2 \times \mathbb{R}^2$; and then the identity is proven in [SS2, Section 2]. Combining all the above we find

$$w(x_1, \dots, x_n) = n^2 \mathcal{F}(\mu_0) + 2n \int \zeta\, d\nu_n + \frac{1}{\pi} W(\nabla H_n, \mathbf{1}_{\mathbb{R}^2}). \tag{2.2}$$

But, changing variables, we have

$$\frac{1}{2}\int_{\mathbb{R}^2\setminus\cup_{i=1}^n B(x_i,\eta)} |\nabla H_n|^2 = \frac{1}{2}\int_{\mathbb{R}^2\setminus\cup_{i=1}^n B(x_i',n\eta)} |\nabla H_n'|^2,$$

and by adding $\pi n \log \eta$ on both sides and letting $\eta \to 0$ we deduce that $W(\nabla H_n, \mathbf{1}_{\mathbb{R}^2}) = W(-\nabla H_n', \mathbf{1}_{\mathbb{R}^2}) - \pi n \log n$. Together with (2.2) this proves the lemma.

2.3 Minimization of W: proof of Theorem 1

By scaling (cf. (1.9)), we reduce to $m = 1$. The result relies on the fact that there exists a minimizing sequence for $\min_{\mathcal{A}_1} W$ consisting of periodic vector-fields:

Proposition 2.1. *For any large enough integer R, there exists a sequence $\{E_R\}_{R\in\mathbb{N}}$ in $\mathcal{A}_1$ such that each E_R is $2R$-periodic (with respect to the x variable) and*

$$\limsup_{R\to\infty} W(E_R) \le \min_{\mathcal{A}_1} W.$$

The proof uses the main results of Section 3 and is postponed to the end of Section 6.

The following proposition could be proven as in [SS1], however we omit the proof here.

Proposition 2.2. $W : L^p_{loc}(\mathbb{R}^2, \mathbb{R}^2) \to \mathbb{R} \cup \{+\infty\}$ *is a Borel function.* $\inf_{\mathcal{A}_1} W$ *is achieved and is finite.*

The result of Theorem 1 will follow from Proposition 2.1 combined with the following

Proposition 2.3 (Minimization in the periodic case)**.** *Let $a_1, \cdots, a_N$ be any points in $[0, N]$ and $E_{\{a_i\}}$ be the corresponding periodic vector field, as in Lemma 1.6. Then*

$$W(E_{\{a_i\}}) \ge W(E_{\mathbb{Z}}) = -\pi \log 2\pi$$

where $E_{\mathbb{Z}}$ is the electric field associated to the perfect lattice $\mathbb{Z}$.

Proof. $W(E_{\mathbb{Z}})$ is immediately computed via (1.10), taking $N = 1$.

Let us now consider arbitrary points $a_1, \dots, a_N$ in $[0, N]$, and assume $a_1 < \cdots < a_N$. Let us also denote $u_{1,i} = a_{i+1} - a_i$, with the convention $a_{N+1} = a_1 + N$. We have $\sum_{i=1}^N u_{1,i} = N$. Similarly, let $u_{p,i} = a_{i+p} - a_i$, with the convention $a_{N+l} = a_l + N$. We have $\sum_{i=1}^N u_{p,i} = pN$. By periodicity of sin, we may view the points a_i as living on the circle $\mathbb{R}/(N\mathbb{Z})$. When adding the terms in $a_i - a_j$ in the sum of (1.10), we can split it according to the difference $p = j - i$ but modulo N. This way, there remains

(2.3)

$$W(E_{\{a_i\}}) = -\frac{\pi}{N}\sum_{i\neq j} \log\left|2\sin\frac{\pi(a_i - a_j)}{N}\right| - \pi\log\frac{2\pi}{N} = -\frac{2\pi}{N}\sum_{p=1}^{[N/2]}\sum_{i=1}^{N}\log\left|2\sin\frac{\pi u_{p,i}}{T}\right| - \pi\log\frac{2\pi}{N},$$

where $[\cdot]$ denotes the integer part. But the function $\log|2\sin x|$ is stricly concave on $[0, \pi]$. It follows that

$$\frac{1}{N}\sum_{i=1}^{N}\log\left|2\sin\frac{\pi u_{p,i}}{N}\right| \le \log\left|2\sin\left(\frac{\pi}{N^2}\sum_{i=1}^{N} u_{p,i}\right)\right| = \log\left|2\sin\frac{p\pi}{N}\right|$$

with equality if and only if all the $u_{p,i}$ are equal. Inserting into (2.3) we obtain

$$W(E_{\{a_i\}}) \geq -2\pi \sum_{p=1}^{[N/2]} \log \left|2\sin\frac{p\pi}{N}\right| - \pi\log\frac{2\pi}{N}. \tag{2.4}$$

On the other hand, if we take for the a_i's the points of the lattice $\mathbb{Z}$ viewing them as N-periodic, we have $u_{p,i} = p$ for all p, i, so if we compute $W(E_{\mathbb{Z}})$ using (2.3), we find

$$W(E_{\mathbb{Z}}) = -\frac{2\pi}{N}\sum_{p=1}^{[N/2]}\sum_{i=1}^{N} \log\left|2\sin\frac{\pi p}{N}\right| - \pi\log\frac{2\pi}{N} = -2\pi\sum_{p=1}^{[N/2]}\log\left|2\sin\frac{\pi p}{N}\right| - \pi\log\frac{2\pi}{N}.$$

This is the right-hand side of (2.4), so (2.4) proves that $W(E_{\{a_i\}}) \geq W(E_{\mathbb{Z}})$ with equality if and only if all the $u_{p,i}$ are equal, which one can easily check implies that $\{a_i\} = \mathbb{Z}$. ☐

Combining with Corollary 2.1, this proves Theorem 1.

3 The main screening result and construction

This section contains the main "upper bound result" relying on explicit constructions, and requiring as a main ingredient the "screening result" mentioned in the introduction. We state the results, whose proofs will occupy the next section.

Proposition 3.1. *Let P be a $T_{\lambda(x)}$-invariant probability measure (see Definition 1.9) on $X = \Sigma \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ with first marginal $dx_{|\Sigma}/|\Sigma|$ and such that for P almost every (x, E) we have $E \in \mathcal{A}_{m_0(x)}$. Then, for any $\eta > 0$, there exists $\delta > 0$ and for any n a subset $A_n \subset \mathbb{R}^n$ such that $|A_n| \geq n!(\delta/n)^n$ and for every sequence $\{\nu_n = \sum_{i=1}^n \delta_{y_i}\}_n$ with $(y_1, \dots, y_n) \in A_n$ the following holds.*

i) We have the upper bound

$$\limsup_{n\to\infty} \frac{1}{n}\left(w_n(y_1, \dots, y_n) - n^2\mathcal{F}(\mu_0) + n\log n\right) \leq \frac{|\Sigma|}{\pi}\int W(E)\, dP(x, E) + \eta. \tag{3.1}$$

ii) There exists $\{E_n\}_n$ in $L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that div $E_n = 2\pi(\nu'_n - m_0{}'\delta_{\mathbb{R}})$ *and such that the image P_n of $dx_{|\Sigma}/|\Sigma|$ by the map $x \mapsto (x, E_n(nx + \cdot))$ is such that*

$$\limsup_{n\to\infty} \textit{dist}\ (P_n, P) \leq \eta, \tag{3.2}$$

where dist is a distance which metrizes the topology of weak convergence on $\mathcal{P}(X)$.

Applying the above proposition with $\eta = 1/k$ we get a subset $A_{n,k}$ in which we choose any n-tuple $(y_{i,k})_{1\leq i\leq n}$. This yields in turn a family $\{P_{n,k}\}$ of probability measures on X. A standard diagonal extraction argument then yields

Corollary 3.2 (Theorem 4, Part B)**.** *Under the same assumptions as Proposition 3.1, there exists a sequence $\{\nu_n = \sum_{i=1}^n \delta_{x_i}\}_n$ and a sequence $\{E_n\}_n$ in $L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that* div $E_n =$ *$2\pi(\nu'_n - m_0{}'\delta_{\mathbb{R}})$ and*

$$\limsup_{n\to\infty} \frac{1}{n}\left(w_n(x_1, \dots, x_n) - n^2\mathcal{F}(\mu_0) + n\log n\right) \leq \frac{|\Sigma|}{\pi}\int W(E)\, dP(x, E). \tag{3.3}$$

Moreover, denoting P_n the image of $dx_{|\Sigma}/|\Sigma|$ by the map $x \mapsto (x, E_n(nx + \cdot))$, we have $P_n \to P$ as $n \to +\infty$.

We may cancel out all leading order terms and rewrite the probability law (1.1) as

$$d\mathbb{P}_n^\beta(x_1,\dots,x_n) = \frac{1}{K_n^\beta} e^{-\frac{n\beta}{2}F_n(\nu)}\, dx_1\dots dx_n \tag{3.4}$$

where

$$K_n^\beta = Z_n^\beta e^{\frac{\beta}{2}(n^2\mathcal{F}(\mu_0)-n\log n)}. \tag{3.5}$$

A consequence of Proposition 3.1 is, recalling (1.27) and (1.35):

Corollary 3.3 (Lower bound part of Theorem 3)**.** *For any $\eta>0$ there exists $C_\eta>0$ such that for any $\beta>0$ we have*

$$\liminf_{n\to+\infty}\frac{\log K_n^\beta}{n}\ge -\frac{\beta}{2}(\alpha+\eta)-C_\eta. \tag{3.6}$$

Proof of the corollary. It is exactly the same as in [SS2] but just letting

$$\sigma_m E(y) := m\, E(my). \tag{3.7}$$

□

We now state our main screening result, on which the proof of Proposition 3.1 is based, and which is the main difference with the two-dimensional situation :

Proposition 3.4. *Let $I_R=[-R/2,R/2]$, let χ_R satisfy (1.7).*

Assume $G\subset\mathcal{A}_1$ is such that there exists $C>0$ such that for any $E\in G$ and writing $\nu=\frac{1}{2\pi}\mathrm{div}\ E+\delta_{\mathbb{R}}$ we have

$$\forall R>1,\ \frac{\nu(I_R)}{R}<C, \tag{3.8}$$

$$\lim_{R\to+\infty}\frac{W(E,\chi_R)}{R}=W(E)<C, \tag{3.9}$$

$$\lim_{y_0\to+\infty}\lim_{R\to+\infty}\fint_{I_R}\int_{|y|>y_0}|E|^2=0, \tag{3.10}$$

and such that moreover all the convergences are uniform w.r.t. $E\in G$.

Then there exists for every $0<\varepsilon<1$, there exists $R_0>0$ such that if $R>R_0$ with $R\in\mathbb{N}$, then for every $E\in G$ there exists a vector field $E_R\in L^p_{loc}(\bar I_R,\mathbb{R}^2)$ such that the following holds:

i) $E_R\cdot\vec\nu=0$ on $\partial\bar I_R$, where $\vec\nu$ denotes the outer unit normal.

ii) There is a discrete subset $\Lambda\subset I_R$ such that

$$\mathrm{div}\ E_R=2\pi\left(\sum_{p\in\Lambda}\delta_p-\delta_{\mathbb{R}}\right)\quad \text{in } \bar I_R.$$

iii) $E_R(z) = E(z)$ *for* $x \in [-R/2 + \varepsilon R, R/2 - \varepsilon R]$.

iv)

$$\frac{W(E_R, \mathbf{1}_{I_R})}{R} \le W(E) + C\varepsilon. \tag{3.11}$$

The assumption (3.10) is a supplementary assumption which is *not* satisfied for all vector-fields in $\mathcal{A}_m$, and which allows to perform the screening. We have not been able to show that screening is always possible without it. However we will see in Section 4 and in particular in Lemma 5.4 that this assumption is satisfied "generically" i.e. for a large set of vector-fields in the support of any invariant probability measure, and this will suffice for our purposes.

4 Proofs of Propositions 3.4 and 3.1

4.1 Preliminaries: a mass displacement result

In this subsection, we state the analogue in 1D of Proposition 4.9 in [SS1], a result we will need later. The point of this result is to say that even though the energy density associated to W: $\frac{1}{2}|E|^2 + \pi \log \eta \sum_p \delta_p$, is not positive or even bounded below, it effectively behaves as if it were, because it can be replaced by an energy-density g which is pointwise bounded below, at the expense of a negligible error. The density g is obtained by displacing the negative part of the energy-density into the positive part. The proof is no different than in [SS1] once the one-dimensional setting has been embedded into the two-dimensional one as stated.

Proposition 4.1. *Assume* (ν, E) *are such that* $\nu = 2\pi \sum_{p\in\Lambda} \delta_p$ *for some finite subset* Λ *of* $\mathbb{R}$, $\operatorname{div} E = 2\pi(\nu - a(x)\delta_{\mathbb{R}})$, *for some* $a \in L^\infty(\mathbb{R})$, *and* $\operatorname{curl} E = 0$. *Then, given* $0 < \rho < \rho_0$, *where* $\rho_0 > 0$ *is universal, there exists a measure density* g *in* $\mathbb{R}^2$ *such that*

i) There exists a family of disjoint closed balls $\mathcal{B}_\rho$ *centered on the real line, covering* $Supp(\nu)$*, such that the sum of the radii of the balls in* $\mathcal{B}_\rho$ *intersected with any segment of* $\mathbb{R}$ *of length* 1 *is bounded by* ρ *and such that*

$$g \ge -C(\|a\|_{L^\infty} + 1) + \frac{1}{4}|E|^2 \mathbf{1}_{\mathbb{R}^2\setminus\mathcal{B}_\rho} \quad \text{in } \mathbb{R}^2, \tag{4.1}$$

where C *depends only on* ρ.

ii)

$$g = \frac{1}{2}|E|^2 \text{ in the complement of } \mathbb{R} \times [-1, 1].$$

iii) For any function χ *compactly supported in* $\mathbb{R}$ *we have, letting* $\bar\chi(x, y) = \chi(x)$,

$$\left| W(E, \chi) - \int \bar\chi \, dg \right| \le CN(\log N + \|a\|_\infty)\|\nabla\chi\|_\infty, \tag{4.2}$$

where $N = \#\{p \in \Lambda \mid B(p, \lambda) \cap Supp(\nabla\bar\chi) \neq \varnothing\}$ *and* λ *depends only on* ρ.

Proposition 4.9 of [SS1] of which the above Proposition is a restatement, was stated for a fixed universal ρ_0, but we may use instead in its proof any $0 < \rho < \rho_0$, which makes the constant C above depend on ρ. Another fact which is true from the proof of [SS1, Proposition 4.9] but not stated in the proposition itself is that in fact $g = \frac{1}{2}|E|^2$ outside $\cup_p B(p,r)$ for some constant $r > 0$ depending only on ρ, and if ρ is taken small enough, then we may take $r = 1$, which yields item ii) of Proposition 4.1.

The next lemma shows that a control on W implies a corresponding control on $\int g$ and of $\int |E|^2$ away from the real axis, growing only like R.

Lemma 4.2. *Assume that $G \subset \mathcal{A}_1$ is such that* (3.8) *and* (3.9) *hold uniformly w.r.t. $E \in G$, for a certain constant C_0. Then for any $E \in G$, writing $\nu = \frac{1}{2\pi}\mathrm{div}\ E + \delta_{\mathbb{R}}$, for every R large enough depending on G, we have*

$$|\nu(I_R) - R| \le C_1 R^{3/4} \log R, \tag{4.3}$$

$$\int_{I_R \times \{|y|>1\}} |E|^2 \le CR\left(W(E) + 1\right), \tag{4.4}$$

and denoting by g the result of applying Proposition 4.1 to E for some fixed value $\rho < 1/8$, we have

$$W(E, \chi_R) - C_1 R^{3/4} \log^2 R \le \int_{\bar{I}_R} dg \le W(E, \chi_{R+1}) + C_1 R^{3/4} \log^2 R, \tag{4.5}$$

where χ_R satisfies (1.7)*, C_1 depends only on G and C is a universal constant.*

Proof. We denote by C_1 any constant depending only on G, and by C any universal constant. From (3.8), (3.9) we have for any $E \in G$ that $\nu(I_R) \le C_1 R$ and $W(E, \chi_R) \le C_1 R$ if R is large enough depending on G. Thus, applying (4.2) we have

$$\left| \int \bar{\chi}_R \, dg \right| \le C_1 R (\log R + 1)$$

which, in view of the fact that $\chi_R = 1$ in I_{R-1} and g is positive outside $\mathbb{R} \times [-1,1]$ and bounded below by a constant otherwise, yields that for every R large enough,

$$\int_{\bar{I}_{R-1}} dg \le C_1 R(\log R + 1). \tag{4.6}$$

This in turn implies — using (4.1) and the fact that $\frac{1}{2}|E|^2 = g$ outside $\mathbb{R} \times [-1,1]$ — the first (unsufficient) control

$$\int_{\{(x,y) | (x,0) \notin \cup \mathcal{B}_\rho\}} |E|^2 \le C_1 R(\log R + 1).$$

Since the sum of the radii of the balls in $\mathcal{B}_\rho$ intersected with any segment of $\mathbb{R}$ of length 1 bounded by $\rho < 1/8$, we deduce by a mean value argument with respect to the variable x that there exists $t \in [0,1]$ such that

$$\int_{\mathbb{R}} \left| E\left(-\frac{R}{2} - t, y\right)\right|^2 + \left| E\left(\frac{R}{2} + t, y\right)\right|^2 dy \le C_1 R(\log R + 1). \tag{4.7}$$

Using now a mean value argument with respect to y, we deduce from (4.6) the existence of $y_R \in [1, 1+\sqrt{R}]$ such that

$$\int_{-\frac{R}{2}-t}^{\frac{R}{2}+t} |E(x,y_R)|^2 + |E(x,-y_R)|^2 \, dx \le C_1\sqrt{R}(\log R + 1). \tag{4.8}$$

Next, we integrate $\operatorname{div} E = 2\pi(\nu - \delta_{\mathbb{R}})$ on the square $[-\frac{R}{2}-t, \frac{R}{2}+t] \times [-y_R, y_R]$. We find using the symmetry property of Corollary 1.5 that

$$\left|\nu(I_{R-\frac{t}{2}}) - R + 2t\right| \le \int_{-y_R}^{y_R} \left|E\left(-\frac{R}{2}-t, y\right)\right| + \left|E\left(\frac{R}{2}+t, y\right)\right| dy + \int_{-\frac{R}{2}-t}^{\frac{R}{2}+t} |E(x,y_R)| \, dx.$$

Using the Cauchy-Schwarz inequality and (4.7)-(4.8), this leads for R large enough to

$$\left|\nu(I_{R-\frac{t}{2}}) - R\right| \le 2 + C_1 R^{3/4}\sqrt{\log R + 1} + C_1\sqrt{y_R}\sqrt{R(\log R + 1)} \le C_1 R^{3/4}(\log R + 1), \tag{4.9}$$

and then — since $\nu(I_R) \ge \nu(I_{R-t/2})$ — to

$$\nu(I_R) - R \ge C_1 R^{3/4} \log R.$$

To prove the same upper bound for $R - \nu(I_R)$ we proceed in the same way, but using a mean value argument to find some $t \in (-1, 0)$ instead of $(0, 1)$ such that (4.7) holds, and then (4.8) also. We deduce as above that (4.9) holds and conclude by noting that since $t \in (-1, 0)$ we have $\nu(I_R) \le \nu(I_{R-t/2})$. This establishes (4.3).

We may bootstrap this information: Indeed (4.3) implies in particular that $\nu(I_R) - \nu(I_{R-1}) \le C_1 R^{3/4} \log R$ and thus we deduce from (4.2) that (4.5) holds:

$$\left|W(E, \chi_R) - \int \bar{\chi}_R \, dg\right| \le C_1 R^{3/4} \log^2 R.$$

Then since $W(E, \chi_R)/R \to W(E)$ as $R \to \infty$ uniformly w.r.t. $E \in G$ and since g is both bounded from below by a universal constant and equal to $\frac{1}{2}|E|^2$ outside $\mathbb{R} \times [-1, 1]$, we deduce (4.4), for R large enough depending on G.

□

Definition 4.3. *Assume $\nu_n = \sum_{i=1}^n \delta_{x_i}$. Letting $\nu_n' = \sum_{i=1}^n \delta_{x_i'}$ be the measure in blown-up coordinates and $E_{\nu_n} = -\nabla H_n'$, where H_n' is defined by* (1.21)*, we denote by g_{ν_n} the result of applying Proposition 4.1 to (ν_n', E_{ν_n}).*

4.2 Some preliminary construction lemmas

The following lemmas serve to estimate the energy of explicit vector fields on boxes, which will be later combined to make test configurations. They are adaptations of [SS1] and rely on elliptic estimates.

Lemma 4.4. *Let $\mathcal{K}$ be the square $[-\frac{L}{2}, \frac{L}{2}]^2$. Let $\varphi \in L^2(\partial\mathcal{K})$ and $a \in L^\infty([-\frac{L}{2}, \frac{L}{2}])$ be such that $\int_{-\frac{L}{2}}^{\frac{L}{2}} a(x)\, dx = -\int_{\partial\mathcal{K}} \varphi$. Then, a_0 being the average of a over $[-\frac{L}{2}, \frac{L}{2}]$, the solution (well defined up to an additive constant) to*

$$\begin{cases} -\Delta u = a\delta_{\mathbb{R}} & \text{in } \mathcal{K} \\ \dfrac{\partial u}{\partial \nu} = \varphi & \text{on } \partial\mathcal{K}. \end{cases} \tag{4.10}$$

satisfies for every $q \in [1,4]$

$$\int_{\mathcal{K}} |\nabla u|^q \le C_q \left(a_0^q L^2 + L^2 \|a - a_0\|^q_{L^\infty([-\frac{L}{2},\frac{L}{2}])} + L^{2-\frac{q}{2}} \|\varphi\|^q_{L^2(\partial\mathcal{K})} \right).$$

Proof. We write the solution u of (4.10) as $u = u_1 + u_2 + u_3$ where

$$\begin{cases} -\Delta u_1 = a_0 \delta_{\mathbb{R}} & \text{in } \mathcal{K} \\ \dfrac{\partial u_1}{\partial \nu} = \bar{\varphi} & \text{on } \partial\mathcal{K} \end{cases} \tag{4.11}$$

where $\bar{\varphi}$ is equal to 0 on the vertical sides of the square and to $\frac{a_0}{2L}$ on both horizontal sides;

$$\begin{cases} -\Delta u_2 = (a - a_0) \delta_{\mathbb{R}} & \text{in } \mathcal{K} \\ \dfrac{\partial u_2}{\partial \nu} = 0 & \text{on } \partial\mathcal{K} \end{cases} \tag{4.12}$$

and

$$\begin{cases} -\Delta u_3 = 0 & \text{in } \mathcal{K} \\ \dfrac{\partial u_3}{\partial \nu} = \varphi - \bar{\varphi} & \text{on } \partial\mathcal{K}. \end{cases}$$

The solution of (4.11) is (up to a constant) $u_1(x,y) = \frac{a_0}{2}|y|$. Hence

$$\int_{\mathcal{K}} |\nabla u_1|^q = \left(\frac{m_0}{2}\right)^q L^2. \tag{4.13}$$

For u_2, we observe that $\|(a-a_0)\delta_{\mathbb{R}}\|_{W^{-1,q}(\mathcal{K})}$ is controlled, for any $q < \infty$, by $\|a-a_0\|_{L^\infty([-\frac{L}{2},\frac{L}{2}])}$. Therefore, using elliptic regularity for (4.12), $\|\nabla u_2\|_{L^q(\mathcal{K})}$ is controlled by $\|a - a_0\|_{L^\infty([-\frac{L}{2},\frac{L}{2}])}$ and a scaling argument shows that for any $q < \infty$

$$\int_{\mathcal{K}} |\nabla u_2|^q \le C_q L^2 \|a - a_0\|^q_{L^\infty([-\frac{L}{2},\frac{L}{2}])}. \tag{4.14}$$

Finally, in the proof of Lemma 4.16 of [SS1] it is shown that for any $q \in [1,4]$

$$\int_{\mathcal{K}} |\nabla u_3|^q \le C_q L^{2-\frac{q}{2}} \|\varphi\|^q_{L^2(\partial\mathcal{K})}. \tag{4.15}$$

Combining (4.13), (4.14) and (4.15), we obtain the result.

□

Lemma 4.5. *Let* m *be a positive constant and let* $\mathcal{K}$ *be a square of center* 0, *and sidelength* $1/m$. *Then the solution to*

$$\begin{cases} -\Delta f = 2\pi(\delta_0 - m\delta_{\mathbb{R}}) & \text{in } \mathcal{K} \\ \dfrac{\partial f}{\partial \nu} = 0 & \text{on } \partial\mathcal{K} \end{cases}$$

satisfies

$$\lim_{\eta \to 0} \left| \int_{\mathcal{K} \setminus B(0,\eta)} |\nabla f|^2 + 2\pi \log \eta \right| = C - \pi \log m \tag{4.16}$$

where C is universal, and for every $1 \leq q < 2$

$$\int_{\mathcal{K}} |\nabla f|^q \leq C_q m^{q-2}, \tag{4.17}$$

where C_q depends only on q.

Proof. By scaling we can reduce to the case of $m = 1$. Then, it suffices to observe that $f(z) = -\log|z| + S(z)$ with $S \in W^{1,\infty}(\mathcal{K})$ and scale back. $\square$

We note that W as defined in (1.6) still makes sense for vector fields satisfying $\mathrm{div}\ E = 2\pi(\sum \delta_{x_i} - m)$ as long as $\mathrm{curl}\, E = 0$ in $\cup B(x_i, \eta_0)$ for some $\eta_0 > 0$. This is the notion we will use repeatedly below.

Lemma 4.6. *Let E_1 and E_2 be two vector fields defined in a rectangle $\mathcal{R}$ of the plane which is symmetric with respect to the real axis, and satisfying*

$$\mathrm{div}\ E_1 = 2\pi(\sum_i \delta_{x_i} - a_1\delta_{\mathbb{R}}) \quad \text{in } \mathcal{R} \tag{4.18}$$

$$\mathrm{div}\ E_2 = a_2\delta_{\mathbb{R}} \quad \text{in } \mathcal{R} \tag{4.19}$$

and $\mathrm{curl}\, E_1$ and $\mathrm{curl}\, E_2$ vanish near the x_i's, for some distinct points $x_i \in \mathbb{R}$ and some bounded functions on the real line, a_1 and a_2. Then, for $q < 2$ and q' its conjuguate exponent, we have $E_1 \in L^q(\mathcal{R})$ and $E_2 \in L^{q'}(\mathcal{R})$ and

$$W(E_1 + E_2, \mathbf{1}_{\mathcal{R}}) \leq W(E_1, \mathbf{1}_{\mathcal{R}}) + \|E_1\|_{L^q(\mathcal{R})}\|E_2\|_{L^{q'}(\mathcal{R})} + \frac{1}{2}\|E_2\|^2_{L^2(\mathcal{R})},$$

where W is still defined as in (1.6)*.*

Proof. We have

$$\int_{\mathcal{R}\setminus\cup_i B(x_i,\eta)} |E_1 + E_2|^2 = \int_{\mathcal{R}\setminus\cup_i B(x_i,\eta)} |E_1|^2 + |E_2|^2 + 2E_1 \cdot E_2.$$

By Hölder's inequality we have

$$\left|\int_{\mathcal{R}\setminus\cup_i B(x_i,\eta)} E_1 \cdot E_2\right| \leq \|E_1\|_{L^q(\mathcal{R})}\|E_2\|_{L^{q'}(\mathcal{R})}$$

The result easily follows.

$\square$

4.3 Proof of the screening result Proposition 3.4

First we note that, in view of (3.10), if we assume $G \subset \mathcal{A}_1$ satisfies the hypothesis of Proposition 3.4 and $0 < \varepsilon < 1$, then there exists $y_0 > 0$ and $R_0 > 0$ such that for all $E \in G$, we have

$$\forall R > R_0, \qquad \int_{I_R \times \{|y| > y_0\}} |E|^2 < \varepsilon^{10} R. \tag{4.20}$$

This motivates the following:

Lemma 4.7. *Let $E \in \mathcal{A}_1$ satisfy* (4.20)*, where $0 < \varepsilon < 1$. Then, denoting $z = (x, y)$, if $|y| > \max(2y_0, R_0)$, we have*

$$|E|^2(z) \le C\frac{\varepsilon^{10}(|x| + |y|)}{|y|^2},$$

where C is universal.

Proof. Each of the coordinates of E is harmonic in the half plane $\mathbb{R}^2_+ = \{y > 0\}$ since $\operatorname{div} E = \operatorname{curl} E = 0$ there. Therefore $|E|^2$ is sub-harmonic. Thus, if $B(z, |y|/2) \subset \mathbb{R}^2_+$ then by the maximum principle we have

$$|E|^2(z) \le \fint_{B(z,|y|/2)} |E|^2.$$

If $y > 2y_0$, then $B(z, |y|/2) \subset [x - \frac{|y|}{2}, x + \frac{|y|}{2}] \times [y_0, +\infty) \subset [-|x| - |y|, |x| + |y|] \times [y_0, +\infty)$. Thus in view of (4.20), if $|x| + |y| > R_0/2$ we have

$$|E|^2(z) \le \frac{8}{\pi}\varepsilon^{10}\frac{|x| + |y|}{y^2},$$

and the result follows, by symmetry with respect to the x-axis. □

Lemma 4.8. *Let G satisfy the assumptions of Proposition 3.4. Then for any $E \in G$, any $0 < \varepsilon < 1/2$ and any R large enough depending on G, ε, we may find $t \in [\frac{R}{2} - \varepsilon R, \frac{R}{2} - \frac{1}{2}\varepsilon R]$ such that*

$$\int_{(\{-t\}\cup\{t\})\times\mathbb{R}} |E|^2 \le C\varepsilon^{-1} \tag{4.21}$$

where C depends only on G, and

$$\lim_{R\to\infty} \frac{1}{2t} W(E, \mathbf{1}_{\bar{I}_{2t}}) = W(E) \tag{4.22}$$

uniformly in G.

Proof. Take $E \in G$ and apply Proposition 4.1 to E for some fixed $0 < \rho < 1/8$. We obtain a density g and balls $\mathcal{B}_\rho$. Now, using (4.5) in Lemma 4.2 together with the bound (3.9), we deduce that if R is large enough depending on G then for any $E \in G$ and denoting g the result of applying Proposition 4.1 to E we have

$$\int_{x=R/2-\varepsilon R}^{R/2-\varepsilon R/2} \int_{\mathbb{R}} (g(x, y) + g(-x, y))\, dx\, dy \le CR + \int_{\bar{I}_R} dg \le CR, \tag{4.23}$$

where C depends only on G and we have used the fact that $g \ge -C$ everywhere and $g \ge 0$ on the set $\{|y| > 1\}$. Then by using the fact that the radii of the balls in $\mathcal{B}_\rho$ which intersect any given interval of length 1 is bounded by $1/8$ we deduce that if R is also large enough depending on ε, the measure of the set A of $x \in [R/2 - \varepsilon R, R/2 - \varepsilon R/2]$ such that $\{x, -x\} \times \mathbb{R}$ does not intersect $\mathcal{B}_\rho$ is bounded below by $\varepsilon R/4$. This and (4.23) implies that the set T of $t \in A$ such that $\int_{\mathbb{R}} (g(t, y) + g(-t, y))\, dy < C/\varepsilon$ has measure at least $\varepsilon R/8$ if C is chosen large enough depending on G, and (4.1) and the fact that $g = \frac{1}{2}|E|^2$ outside $\mathbb{R} \times [-1, 1]$ imply that (4.21) holds for $t \in T$. Thus

$$|\{t \in [R/2 - \varepsilon R, R/2 - \varepsilon R/2] \mid (4.21) \text{ holds}\}| \geq \frac{\varepsilon R}{8}. \tag{4.24}$$

For (4.22) we argue as in [SS1], Lemma 4.14. We let $\chi : [0, +\infty) \to \mathbb{R}$ be a monotonic function with compact support and let $\bar{\chi}(x, y) = \chi(|x|)$. First we note that for any Radon measure μ in $\mathbb{R}^2$ we have

$$\int \bar{\chi}\, d\mu = -\int_{t=0}^{+\infty} \chi'(t)\mu(\bar{I}_{2t})\, dt = -2\int_{u=0}^{+\infty} \chi'(u/2)\mu(\bar{I}_u)\, du.$$

This implies straightforwardly using the definition of $W(E, \chi)$ that

$$2\int_{u=0}^{+\infty} \left(W(E, \mathbf{1}_{\bar{I}_u}) - g(\bar{I}_u)\right) \chi'(u/2)\, dt = \int \bar{\chi}\, dg - W(E, \chi).$$

On the other hand, by (4.2) and applying Lemma 4.2, (4.3), if χ' is supported in $[x, y] \subset [R/2, R]$, then the right-hand side is bounded by $C(|x - y| + R^{3/4}\log^2 R)\|\chi'\|_\infty$ for any R large enough depending on G. Given now any $\rho : \mathbb{R}_+ \to \mathbb{R}$ supported in $[x, y] \subset [R/2, R]$ we may consider the positive and negative parts ρ_+ and ρ_-, and and their primitives χ_+ and χ_- with compact support, which are monotonic. Applying the above to χ_+ and χ_- we find

$$\int_{u=0}^{+\infty} \left(W(E, \mathbf{1}_{\bar{I}_u}) - g(\bar{I}_u)\right) \rho(u)\, du \leq C(|x - y| + R^{3/4}\log^2 R)\|\rho\|_\infty.$$

Since this is true for any ρ supported in $[R - 2\varepsilon R, R - \varepsilon R]$, it follows by duality that

$$\int_x^y |W(E, \mathbf{1}_{\bar{I}_u}) - g(\bar{I}_u)|\, du \leq C(|x - y| + R^{3/4}\log^2 R). \tag{4.25}$$

Then we divide $[R - 2\varepsilon R, R - \varepsilon R]$ into, say, $[\sqrt{R}]$ intervals $I_1, \dots, I_{[\sqrt{R}]}$ so that their length is equivalent to $\varepsilon\sqrt{R}$ for large R. Then, for large enough R, on each such interval (4.25) implies that

$$\int_{I_k} |W(E, \mathbf{1}_{\bar{I}_u}) - g(\bar{I}_u)|\, du \leq CR^{3/4}\log^2 R.$$

Therefore the set of $u \in I_k$ such that $|W(E, \mathbf{1}_{\bar{I}_u}) - g(\bar{I}_u)| \leq 8CR^{3/4}\log^2 R$ has measure at least $7|I_k|/8$ and since this is true on each interval, and changing variables, the set of $t \in [R/2 - \varepsilon R, R/2 - \varepsilon R/2]$ such that $|W(E, \mathbf{1}_{\bar{I}_{2t}}) - g(\bar{I}_t)| \leq 8CR^{3/4}\log^2 R$ has measure at least $7\varepsilon R/16$. Together with (4.24), this implies the existence of $t \in [R/2 - \varepsilon R, R/2 - \varepsilon R/2]$ such that both (4.21) and (4.22) hold.

□

We now prove Proposition 3.4. Let G satisfy its hypothesis and choose $0 < \varepsilon < 1$, and $E \in G$. Applying Lemma 4.8 we find that if R is large enough depending on G, ε, then there exists $t \in [R/2 - \varepsilon R, R/2 - \varepsilon R/2]$ such that (4.21) and (4.22) hold. For any such *integer* $R \in \mathbb{N}$ we may also choose y_R such that

$$\varepsilon^3 R < y_R < \varepsilon^{5/2} R. \tag{4.26}$$

Finally we choose $s > t$ such that $s - t \in [y_R, y_R + 1]$ and $\frac{R}{2} - s \in \mathbb{N}$, and start constructing the vector field E_R.

- *Step 1: splitting the strip.* We split the strip $I_R = [-\frac{R}{2}, \frac{R}{2}] \times \mathbb{R}$ into several rectangles and strips (see figure below): let

$$\begin{aligned}
D_0 &= [-t, t] \times [-y_R, y_R] \\
D_+ &= [t, s] \times [-y_R, y_R] \\
D_- &= [-s, -t] \times [-y_R, y_R] \\
D_e^+ &= [s, R/2] \times \mathbb{R} \\
D_e^- &= [-R/2, -s] \times \mathbb{R} \\
D_1 &= [-s, s] \times ([y_R, y_R + R] \cup (-R - y_R, -y_R]) \\
D_\infty &= [-s, s] \times ([R + y_R, +\infty) \cup (-\infty, -R - y_R]) \,.
\end{aligned}$$



First we let $E_R = E$ in D_0, $E_R = 0$ in D_∞ and below we are going to define E_R on each of the other sets.

Recall that from Corollary 1.5 we have that $E(x, y)$ is the reflection of $E(x, -y)$ with respect to the line $\{y = 0\}$. We denote by φ_+ the trace $E \cdot \vec{\nu}$ on the right-hand side of D_0 where $\vec{\nu}$ is the outward-pointing normal to D_0, φ_- the same on the left-hand side, φ_h the trace on the upper side of D_0 (which by symmetry of the problem with respect to the real axis is equal to that on the lower side). From (4.21) and the Cauchy-Schwarz inequality, we have

$$\int |\varphi_-|^2 + \int |\varphi_+|^2 \leq \frac{C}{\varepsilon} \qquad \int |\varphi_-| + \int |\varphi_+| \leq C\sqrt{\frac{y_R}{\varepsilon}}. \tag{4.27}$$

and from Lemma 4.7 and (4.26) we have

$$\int_{[-t,t]} |\varphi_h|^2 \leq \frac{C\varepsilon^{10} R(R + y_R)}{y_R^2} < C\varepsilon^4 \qquad \int_{[-t,t]} |\varphi_h| \leq C\varepsilon^2\sqrt{R}. \tag{4.28}$$

In addition, integrating the relation div $E = 2\pi(\sum_{p\in\Lambda}\delta_p - \delta_{\mathbb{R}})$ over D_0 gives

$$\frac{1}{2\pi}\left(\int \varphi_+ + \int \varphi_- + 2\int \varphi_h\right) = \nu([-t,t]) - 2t. \tag{4.29}$$

- *Step 2: defining E_R in D_+.* We first define $\varphi_0 := E_R \cdot \vec{\nu}$ on the boundary ∂D_+, where $\vec{\nu}$ is the outward normal to D_+. For a certain constant φ_h^+ to be chosen later we let

$$\varphi_0 = \begin{cases} -\varphi_+ & \text{on } \partial D_+ \cap \partial D_0, \\ 0 & \text{on } \partial D_+ \cap \{x = s\}, \\ \varphi_h^+ & \text{on } \partial D_+ \cap \{y = \pm y_R\}. \end{cases} \tag{4.30}$$

Then, inside D_+, we let $E_R = E_1 + E_2$, where

$$\begin{cases} \text{div } E_1 = 2\pi\left(\displaystyle\sum_{i=1}^{n_+} \delta_{x_i} - m_+\delta_{\mathbb{R}}\right) & \text{in } \mathcal{D}_+ \\ E_1 \cdot \nu = 0 & \text{on } \partial D_+. \end{cases}, \tag{4.31}$$

$$\begin{cases} \text{div } E_2 = 2\pi(m_+ - 1)\delta_{\mathbb{R}} & \text{in } \mathcal{D}_+ \\ E_2 \cdot \nu = \varphi_0 & \text{on } \partial D_+. \end{cases} \tag{4.32}$$

Here, n_+ is an integer and m_+ a real number which are defined by

$$n_+ = \left[(s-t) - \frac{1}{2\pi}\left(\int \varphi_+ + \int \varphi_h\right)\right], \quad m_+ = \frac{n_+}{s-t}, \tag{4.33}$$

and for $1 \le i \le n_+$ we have let

$$x_i = t + \frac{s-t}{n_+}\left(i + \frac{1}{2}\right).$$

Note that the above equations do not yield a uniquely defined E_1 and E_2. For (4.31) to make sense we need $n_+ \ge 0$ while for (4.32) to have a solution we need to have

$$2\pi\left(n_+ - (s-t)\right) = \int \varphi_0 = 2(s-t)\varphi_h^+ - \int \varphi_+, \tag{4.34}$$

which we take as the definition of φ_h^+. The fact that $n_+ \ge 0$ follows for R large enough depending on ε from the fact that $s - t \ge \varepsilon^3 R$ and (4.27), (4.28).

- *Step 3: Estimating the energy of E_R in D_+, D_-.* To compute the renormalized energy $W(E_R, \mathbf{1}_{D_+})$ we need to define E_1 and E_2 more precisely. For E_1 let us consider n_+ identical squares $\{K_i\}_{i=1}^{n_+}$ with sidelength $\frac{s-t}{n_+} = \frac{1}{m_+}$, sides parallel to the axes and such that K_i is centered at x_i. We define E_1 restricted to K_i by applying Lemma 4.5 with $m = m_+$ and taking $E_1 = -\nabla f$, while outside $\cup_i K_i$ we let $E_1 = 0$. Since from Lemma 4.5 we have $E_{1|K_i} \cdot \vec{\nu} = 0$ on ∂K_i, it holds that div $E_1 = \sum_i$ div $E_{1|K_i}$ and therefore (4.31) is satisfied by

E_1. On the other hand, still from Lemma 4.5 we obtain by summing the bounds (4.16) and (4.17) on the n_+ rectangles

$$\lim_{\eta\to 0}\left|\int_{D_+\setminus\cup_i B(x_i,\eta)} |E_1|^2 + 2\pi\log\eta\right| \le n_+\left(C - \pi\log m_+\right), \tag{4.35}$$

and

$$\forall 1 < q < 2, \quad \int_{D_+} |E_1|^q \le C_q n_+. \tag{4.36}$$

We define E_2 by applying Lemma 4.4 in D_+, hence with $L = s-t$, with the boundary data φ_0 and constant weight $m = 2\pi(m_+ - 1)$. From (4.30) and (4.34) the hypothesis $\int m(x)\,dx = -\int_{\partial D_+}\varphi$ is satisfied and applying the lemma yields

$$\forall 2 \le q < 4, \quad \int_{D_+} |E_2|^q \le C_q\left(|m_+ - 1|^q (s-t)^2 + (s-t)^{2-\frac{q}{2}}\|\varphi_0\|^q_{L^2(\partial D_+)}\right). \tag{4.37}$$

Using Lemma 4.6 we have, recalling that $E_R := E_1 + E_2$ in D_+ and using (4.35), (4.36), (4.37):

$$\begin{aligned} W(E_R, \mathbf{1}_{D_+}) \le Cn_+ + C_q {n_+}^{1/q}\left(|m_+ - 1|(s-t)^{2/q'} + (s-t)^{2/q' - 1/2}\|\varphi_0\|_{L^2(\partial D_+)}\right) \\ + C\left(|m_+ - 1|^2 (s-t)^2 + (s-t)\|\varphi_0\|^2_{L^2(\partial D_+)}\right), \end{aligned} \tag{4.38}$$

for any $1 < q < 2$ such that the conjugate exponent q' is less then 4. Now, from (4.33), we have using (4.27), (4.28), (4.26) and the fact that $y_R \le s - t \le y_R + 1$ we deduce that

$$|n_+ - (s-t)| \le C\left(\varepsilon^2\sqrt{R} + \sqrt{\frac{y_R}{\varepsilon}}\right), \quad |m_+ - 1| \le C\left(\frac{1}{\varepsilon\sqrt{R}} + \frac{1}{\sqrt{\varepsilon y_R}}\right), \tag{4.39}$$

and thus for R large enough depending on ε, since $s - t \simeq y_R$ for large R and using (4.26) again as well as $\varepsilon < 1$,

$$n_+ \le C\varepsilon^{5/2} R, \quad |m_+ - 1| \le \frac{C}{\varepsilon^2\sqrt{R}}. \tag{4.40}$$

Moreover, from (4.33), (4.34) and (4.28) we find

$$|\varphi_h^+| \le \frac{1}{2(s-t)}\left|2\pi + \int\varphi_h\right| \le C\frac{\varepsilon^2\sqrt{R}}{y_R} \le \frac{C}{\varepsilon\sqrt{R}}. \tag{4.41}$$

Then, in view of (4.30), (4.27),

$$\|\varphi_0\|^2_{L^2(\partial D_+)} \le \frac{C}{\varepsilon}.$$

Now we fix for instance $q = 3/2$, so that $q' = 3$ and combining the above with (4.38), (4.40) we find that for R large enough depending on ε, and denoting by C_ε a positive constant depending on ε but independent of R,

$$W(E_R, \mathbf{1}_{D_+}) \le C\varepsilon^{5/2} R + C_\varepsilon R^{2/3} \times R^{2/3 - 1/2} + C\varepsilon^{3/2} R.$$

Thus for R large enough depending on ε we find that

$$W(E_R,\mathbf{1}_{D_+})\le C\varepsilon^{3/2}R. \tag{4.42}$$

An almost symmetric computation yields the same bound for $W(E_R,\mathbf{1}_{D_-})$. It suffices to let

$$n_-=2(s-t)-\frac{1}{2\pi}\left(\int\varphi_+ +\int\varphi_- +2\int\varphi_h\right)-n_+, \tag{4.43}$$

and carry on the proof with minuses instead of pluses. The fact that n_- is an integer follows from the identity (4.29), the fact that $2s$ is an integer and the fact that $\nu([-t,t])\in\mathbb{N}$. Moreover the definition of n_- implies that

$$n_-=\left[(s-t)-\frac{1}{2\pi}\left(\int\varphi_- +\int\varphi_h\right)\right],\quad\text{or}\quad n_-=\left[(s-t)-\frac{1}{2\pi}\left(\int\varphi_- +\int\varphi_h\right)\right]+1,$$

hence n_- is positive if R is large enough and (4.39) holds for n_- as well. The rest of the proof is unchanged.

- *Step 4: defining E_R over D_1.* We need only consider the intersection of D_1 with the upper half-plane (and then extend by reflection). We let φ_0 be equal to $-\varphi_h^-$, $-\varphi_h$, $-\varphi_h^+$, respectively, on the intersection of ∂D_1 with ∂D_-, ∂D_0, ∂D_+, respectively. On the remaining three sides of ∂D_1 we let $\varphi_0=0$. From (4.34) and its equivalent for n_- and the fact that $n_\pm=(s-t)m_\pm$ we have

$$-\int\varphi_0=\pi(n_+ +n_- -2(s-t))+\frac{1}{2}\int\varphi_+ +\frac{1}{2}\int\varphi_- +\int\varphi_h,$$

and then (4.33), (4.43) imply that the integral of φ_0 is zero.

Thus there exists a harmonic function u in D_1 with normal derivative φ_0 on ∂D_1, we let $E_R=\nabla u$ on D_1. Using (4.41) — which holds for φ_h^- as well — and (4.28) we have $C\varepsilon^4+C\varepsilon^4R/y_R$ hence

$$\|\varphi_0\|^2_{L^2(\partial D_1)}\le C\left(\varepsilon^4+\varepsilon^4R/y_R\right)\le C\varepsilon. \tag{4.44}$$

Then standard elliptic estimates yield as in Lemma 4.4 that

$$\int_{D_1}|E_R|^2\le CR\|\varphi_0\|^2_{L^2}\le CR\varepsilon, \tag{4.45}$$

where we have concluded by (4.26).

- *Step 5: defining E_R over D_e^+.* The construction will be entirely parallel in D_e^-. We note that D_e^+ is an infinite strip of width $R/2-s$ and we have chosen s so that this quantity is an integer. We can thus split this strip into exactly $R/2-s$ strips of width 1. On each of these strips we define E_R to be equal to 0 for $|y|\ge\frac{1}{2}$ and for $|y|\le\frac{1}{2}$ (i.e. in a square of sidelength 1) we choose it to be ∇f where f is given by Lemma 4.5 applied with $m=1$. Since $E_R\cdot\vec{\nu}=0$ on the boundary of each of these squares, no divergence is created at the interfaces, and the resulting E_R satisfies $\operatorname{div}E_R=2\pi(\sum_{p\in\Lambda}\delta_p-\delta_{\mathbb{R}})$. In addition in view of (4.17) the cost in energy is equal to a constant times the number of strips, that is

$$W(E_R,\mathbf{1}_{D_e})\le C|R/2-s|\le C\varepsilon R. \tag{4.46}$$

- *Conclusion.* We have now defined E_R over the whole strip $\bar{I}_R$. It satisfies items ii) and iii). The main point is again that as long as $E \cdot \vec{\nu}$ is continuous across an interface it creates no singular divergence there. Combining (4.46) with (4.22), (4.42) and (4.45), E_R also satisfies (3.11). This concludes the proof of Proposition 3.4.

5 Proof of Proposition 3.1

The construction consists of the following. First we select a finite set of vector fields $J_1, \dots, J_N$ (N will depend on ε) which will represent the probability $P(x, E)$ with respect to its E-dependence, and whose renormalized energies are well-controlled. Since P is $T_{\lambda(x)}$-invariant, we need it to be well-approximated by measures supported on the orbits of the J_i's under translations. It is also during this approximation process that we manage to select the J_i's as belonging to a part of the support of P of almost full measure for which the extra assumption (3.10) holds and the screening can be made. Secondly, we work in blown-up coordinates and split the region Σ' (of order n size) into many intervals, and then select the proportion of the intervals that corresponds to the weight that the orbit of each J_i carries in the approximation of P. In these rectangles we paste a (translated) copy of J_i at the appropriate scale (approximating the density ${m_0}'$ by a piecewise constant one and controlling errors).

To conclude the proof of Proposition 3.1, we collect all of the estimates on the constructed vector field to show that its energy w_n is bounded above in terms of $\int W \, dP$ and that the probability measures associated to the construction have remained close to P.

5.1 Abstract preliminaries

We repeat here the definitions of distances that we used in [SS2]. First we choose distances which metrize the topologies of $L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ and $\mathcal{B}(X)$, the set of finite Borel measures on $X = \Sigma \times L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$. For $E_1, E_2 \in L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ we let

$$d_p(E_1, E_2) = \sum_{k=1}^{\infty} 2^{-k} \frac{\|E_1 - E_2\|_{L^p(B(0,k))}}{1 + \|E_1 - E_2\|_{L^p(B(0,k))}},$$

and on X we use the product of the Euclidean distance on Σ and d_p, which we denote d_X. On $\mathcal{B}(X)$ we define a distance by choosing a sequence of bounded continuous functions $\{\varphi_k\}_k$ which is dense in $C_b(X)$ and we let, for any $\mu_1, \mu_2 \in \mathcal{B}(X)$,

$$d_{\mathcal{B}}(\mu_1, \mu_2) = \sum_{k=1}^{\infty} 2^{-k} \frac{|\langle \varphi_k, \mu_1 - \mu_2 \rangle|}{1 + |\langle \varphi_k, \mu_1 - \mu_2 \rangle|},$$

where we have used the notation $\langle \varphi, \mu \rangle = \int \varphi \, d\mu$.

We will use the following general facts, whose proofs are in [SS2, Sec. 7.1].

Lemma 5.1. *For any $\varepsilon > 0$ there exists $\eta_0 > 0$ such that if $P, Q \in \mathcal{B}(X)$ and $\|P - Q\| < \eta_0$, then $d(P, Q) < \varepsilon$. Here $\|P - Q\|$ denotes the total variation of the signed measure $P - Q$, i.e. the supremum of $\langle \varphi, P - Q \rangle$ over measurable functions φ such that $|\varphi| \le 1$.*

In particular, if $P = \sum_{i=1}^{\infty} \alpha_i \delta_{x_i}$ and $Q = \sum_{i=1}^{\infty} \beta_i \delta_{x_i}$ with $\sum_i |\alpha_i - \beta_i| < \eta_0$, then $d_{\mathcal{B}}(P, Q) < \varepsilon$.

Lemma 5.2. *Let $K \subset X$ be compact. For any $\varepsilon > 0$ there exists $\eta_1 > 0$ such that if $x \in K, y \in X$ and $d_X(x,y) < \eta_1$ then $d_{\mathcal{B}}(\delta_x, \delta_y) < \varepsilon$.*

Lemma 5.3. *Let $0 < \varepsilon < 1$. If μ is a probability measure on a set A and $f, g : A \to X$ are measurable and such that $d_{\mathcal{B}}(\delta_{f(x)}, \delta_{g(x)}) < \varepsilon$ for every $x \in A$, then*

$$d_{\mathcal{B}}(f\#\mu, g\#\mu) < C\varepsilon(|\log \varepsilon| + 1)$$

where $\#$ denotes the push-forward of a measure.

The next lemma shows how to, given a translation-invariant probability measure $\tilde{P}$ on $L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$, select a good subset G_ε and vector fields J_i of $L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ to approximate it. It is essentially borrowed from [SS2] except it contains in addition the argument that ensures that we may choose G_ε to satisfy the assumption (3.10) needed for the screening.

Lemma 5.4. *Let $\tilde{P}$ be a translation invariant measure on X such that, $\tilde{P}$-a.e., E is in $\mathcal{A}_1$ and satifies $W(E) < +\infty$. Then, for any $\varepsilon > 0$ there exists a compact $G_\varepsilon \subset L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that*

i) Letting $0 < \eta_0$ be as in Lemma 5.1 we have

$$\tilde{P}(\Sigma \times {G_\varepsilon}^c) < \min({\eta_0}^2, \eta_0 \varepsilon). \tag{5.1}$$

ii) The convergence (1.8) is uniform with respect to $E \in G_\varepsilon$.

iii) Writing $\mathrm{div}\ E = 2\pi(\nu_E - 1)$, both $W(E)$ and $\nu_E(I_R)/R$ are bounded uniformly with respect to $E \in G_\varepsilon$ and $R > 1$.

iv) Uniformly with respect to $E \in G_\varepsilon$ we have

$$\lim_{y_0 \to +\infty} \lim_{R \to +\infty} \fint_{I_R} \int_{|y| > y_0} |E|^2 = 0 \tag{5.2}$$

Moreover, (5.1) implies that for any $R_\varepsilon > 1$ there exists a compact subset $H_\varepsilon \subset G_\varepsilon$ such that

v) For every $E \in H_\varepsilon$, there exists $\Gamma(E) \subset I_{\overline{m}R_\varepsilon}$ such that

$$|\Gamma(E)| < CR_\varepsilon \eta_0 \text{ and } \lambda \notin \Gamma(E) \implies \theta_\lambda E \in G_\varepsilon. \tag{5.3}$$

vi) We have

$$d_{\mathcal{B}}(\bar{P}, P') < C\varepsilon(|\log \varepsilon| + 1), \quad \text{where} \tag{5.4}$$

$$P' = \int_{\Sigma \times H_\varepsilon} \frac{1}{m_0(x)|I_{R_\varepsilon}|} \int_{m_0(x) I_{R_\varepsilon} \setminus \Gamma(E)} \delta_x \otimes \delta_{\sigma_{m_0(x)} \theta_\lambda E} \, d\lambda \, d\tilde{P}(x, E)$$

$$\bar{P} = \int_{\Sigma \times L^p_{\mathrm{loc}}(\mathbb{R}^2, \mathbb{R}^2)} \delta_x \otimes \delta_{\sigma_{m_0(x)} E} \, d\tilde{P}(x, E).$$

Finally, there exists a partition of H_ε *into* $\cup_{i=1}^{N_\varepsilon} H_\varepsilon^i$ *satisfying* diam $(H_\varepsilon^i) < \eta_3$*, where* η_3 *is such that*

$$E \in H_\varepsilon,\ d_p(E,E') < \eta_3,\ m \in (0,\overline{m}],\ \lambda \in \overline{m} I_{R_\varepsilon} \backslash \Gamma(E) \implies d_{\mathcal{B}}(\delta_{\sigma_m \theta_\lambda E}, \delta_{\sigma_m \theta_\lambda E'}) < \varepsilon; \tag{5.5}$$

and there exists for all i*,* $E_i \in H_\varepsilon^i$ *such that*

$$W(E_i) < \inf_{H_\varepsilon^i} W + \varepsilon. \tag{5.6}$$

Proof. The lemma is almost identical to Lemma 7.6 in [SS2], except for item iv). The proof in [SS2] is as follows: First one proves that there exists G_ε satisfying items ii) and iii) with $\tilde{P}(\Sigma \times {G_\varepsilon}^c)$ arbitrarily small, in particular one can choose it so that (5.1) is satisfied. Then one deduces from (5.1) the existence, for any $R_\varepsilon > 1$, of a compact subset $H_\varepsilon \subset G_\varepsilon$ satisfying the remaining properties. The only difference here is that we must check that there exists G_ε with $\tilde{P}(\Sigma \times {G_\varepsilon}^c)$ arbitrarily small satisfying not only items ii) and iii), but iv) as well. Then, the proof of the existence $H_\varepsilon \subset G_\varepsilon$ satisfying the remaining properties is exactly as in [SS2].

Of course, by intersecting sets, it is equivalent to prove that ii), iii), and iv) can be satisfied *simultaneously* or *separately*, on a set of measure arbitrarily close to full. The proof in [SS2] shows that this is possible for ii) or iii), it remains to check it for iv). For this we consider $G_n = \{E \mid W(E) < n\}$. Then G_n is a translation-invariant set since W is a translation-invariant function, and therefore by the multiparameter ergodic theorem (as in [Be]), and since $\tilde{P}$ is translation-invariant, we have

$$\int_{\Sigma \times G_n} \left(\int_{[-1,1]\times\{|y|>y_0\}} |E|^2 \right) d\tilde{P}(x,E) = \int_{\Sigma \times G_n} \left(\lim_{R\to\infty} \frac{1}{R} \int_{\mathbb{R}\times\{|y|>y_0\}} \chi_R |E|^2 \right) d\tilde{P}(x,E), \tag{5.7}$$

where $\chi_R = \mathbf{1}_{I_R} * \mathbf{1}_{[-1,1]}$. Then, using Lemma 4.2 and using the fact that the g there was defined in Proposition 4.1 hence is equal to $\frac{1}{2}|E|^2$ on $\mathbb{R} \times \{|y| > 1\}$ we deduce from (4.5) and the fact that g is bounded below by a constant independent of E that

$$\lim_{R\to\infty} \frac{1}{R} \int_{\mathbb{R}\times\{|y|>y_0\}} \chi_R |E|^2 \le C(1 + W(E)) \le Cn$$

holds for every $E \in G_n$ with $n \ge 1$.

It follows that for every fixed $n \ge 1$ the family of functions

$$\left\{ \varphi_{y_0} : (x,E) \mapsto \int_{[-1,1]\times\{|y|>y_0\}} |E|^2 \right\}_{y_0>1}$$

decreases to 0 on $\Sigma \times G_n$ as $y_0 \to +\infty$, and is dominated by the bounded, hence $\tilde{P}$-integrable, function φ_1. Lebesgue's theorem then implies that their integrals on $\Sigma \times G_n$ converge to 0, hence in view of (5.7) that

$$\int_{\Sigma \times G_n} \left(\lim_{R\to\infty} \frac{1}{R} \int_{\mathbb{R}\times\{|y|>y_0\}} \chi_R |E|^2 \right) d\tilde{P}(x,E),$$

tends to 0 as $y_0 \to +\infty$. Fatou's lemma then implies that (5.2) holds for $\tilde{P}$-almost every $(x,E) \in \Sigma \times G_n$.

Since $W(E) < +\infty$ holds for $\tilde{P}$-a.e. (x,E), we know that $\tilde{P}(\Sigma \times G_n) \to 1$ as $n \to +\infty$ therefore the measure of $\Sigma \times G_n$ can be made arbitrarily close to 1, and then Egoroff's theorem implies that by restricting G_n we can in addition require the convergence in (5.2) to be uniform. □

5.2 Construction

In what follows $\Sigma' = n\Sigma$, ${m_0}'(x) = m_0(x/n)$: we work in blown-up coordinates. In view of assumption (1.16), we may assume without loss of generality that Σ is made of one closed interval $[a,b]$ (it is then immediate to generalize the construction to the case of a finite union of intervals). In that case $\Sigma' = [na, nb]$. Let $\underline{m} > 0$ be a small parameter. For any integer n we choose real numbers a_n and b_n (depending on $\underline{m}$) as follows: Let a_n be the smallest number and b_n the largest such that

$$a_n \geq na + \frac{n\underline{m}}{\gamma^2} \qquad b_n \leq nb - \frac{n\underline{m}}{\gamma^2} \tag{5.8}$$

$$\int_{na}^{a_n} {m_0}'(x)\,dx \in \mathbb{N} \tag{5.9}$$

$$\int_{b_n}^{nb} {m_0}'(x)\,dx \in \mathbb{N} \tag{5.10}$$

$$\int_{a_n}^{b_n} {m_0}'(x)\,dx \in q_\varepsilon\mathbb{N} \tag{5.11}$$

where q_ε is an integer, to be chosen later, and γ is the constant in (1.17). By (5.8) and assumption (1.16), we are sure to have ${m_0}' \geq \underline{m}$ in $\Sigma'_{\underline{m}} := [a_n, b_n]$. This fact also ensures that

$$|a_n - na| \leq \frac{n\underline{m}}{\gamma^2} + \frac{1}{\underline{m}} \qquad |b_n - nb| \leq \frac{n\underline{m}}{\gamma^2} + \frac{q_\varepsilon}{\underline{m}}. \tag{5.12}$$

We also denote $\Sigma_{\overline{m}} := \frac{1}{n}\Sigma'_{\overline{m}}$.

Let P be a probability on $\Sigma \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ which is as in the statement of Proposition 3.1. Our goal is to construct a vector field E_n whose W energy is close to $\int W\,dP$ and such that the associated P_n (defined as the push-forward of the normalized Lebesgue measure on Σ by $x \mapsto (x, E(nx + \cdot))$) well approximates P, a probability measure as in the statement of Proposition 3.1.

In $[na, a_n]$ and $[b_n, nb]$, we approximate ${m_0}'(x)\,dx$ by a sum of Dirac masses at points appropriately spaced, and build an associated E_n, whose contribution to the energy will shown to be negligible as $\overline{m} \to 0$. We leave this part for the end.

For now we turn to $[a_n, b_n]$, where we will do a more sophisticated construction, approaching P via Lemma 5.4 and using Proposition 3.4. The idea of the construction is to split the interval $[a_n, b_n]$ into intervals of width $\sim q_\varepsilon R_\varepsilon$, where q_ε is an integer and R_ε a number, both chosen large enough, and then paste in each of these intervals a large number of copies of the (rescaled) truncations of the J_i's provided by Proposition 3.4, in a proportion following that of P.

-Step 1: Reduction to a density bounded below. We have

$$P = \int \delta_x \otimes \delta_{\sigma_{m_0(x)}E}\,dQ(x,E), \quad \text{where} \quad Q = \int \delta_x \otimes \delta_{\sigma_{1/m_0(x)}E}\,dP(x,E).$$

Moreover, since the first marginal of P is the normalized Lebesgue measure on Σ and since $|\Sigma_{\underline{m}}| \simeq |\Sigma|$ as $\underline{m} \to 0$, we have

$$\lim_{\underline{m}\to 0} d_{\mathcal{B}}(P, \bar{P}) = 0, \tag{5.13}$$

where $\bar P$ is defined by

$$\bar P = \int \delta_x \otimes \delta_{\sigma_{m_0(x)}E}\, d\tilde P(x,E), \quad \text{with} \quad \tilde P = \int_{\Sigma_{\underline{m}} \times L^p_{\mathrm{loc}}(\mathbb{R}^2,\mathbb{R}^2)} \delta_x \otimes \delta_{\sigma_{1/m_0(x)}E}\, dP(x,E). \tag{5.14}$$

Clearly $\tilde P$ is is $T_{\lambda(x)}$-invariant since P is, and in particular it is translation-invariant. In addition, for $\tilde P$-a.e. (x,E), we have $m_0(x) \in [\underline{m}, \overline{m}]$, a situation similar to [SS2] where the density was assumed to be bounded below.

-Step 2: Choice of the parameters. Let $0 < \varepsilon < 1$. We apply Lemma 5.4, which provides us with a compact set $G_\varepsilon \subset L^p_{\mathrm{loc}}(\mathbb{R}^2,\mathbb{R}^2)$. We may then apply Proposition 3.4 with the same ε to G_ε. It provides us with $R_0 > 0$ such that for any integer $R > R_0$, and roughly speaking, any $E \in G_\varepsilon$ may be truncated on $\bar I_R$ with an ε-cost in energy. Applying Lemma 5.2 on the compact set $\{\sigma_m E : m \in [\underline{m},\overline{m}], E \in G_\varepsilon\}$, there exists $\eta_1 > 0$ such that

$$m \in [\underline{m},\overline{m}],\ E \in G_\varepsilon, E' \in L^p_{\mathrm{loc}}(\mathbb{R}^2,\mathbb{R}^2) \text{ and } d_p(E,E') < \eta_1 \implies d_{\mathcal{B}}(\delta_{\sigma_m E}, \delta_{\sigma_m E'}) < \varepsilon. \tag{5.15}$$

Then we define R_ε to be such that $\underline{m} R_\varepsilon > R_0$ and such that for any $E, E' \in L^p_{\mathrm{loc}}(\mathbb{R}^2,\mathbb{R}^2)$,

$$E = E' \text{ on } \bar I_{\underline{m}\varepsilon R_\varepsilon} \quad \Longrightarrow \quad d_p(E,E') < \eta_1. \tag{5.16}$$

Going back to Lemma 5.4, we deduce the existence of $H_\varepsilon \subset G_\varepsilon$, of $N_\varepsilon \in \mathbb{N}$ and of $\{E_i\}_{1 \le i \le N_\varepsilon}$ satisfying (5.3), (5.4), (5.5) and (5.6).

Finally, we choose $q_\varepsilon \in \mathbb{N}$ sufficiently large so that

$$\frac{N_\varepsilon}{q_\varepsilon} < \eta_0, \qquad \frac{N_\varepsilon}{{q_\varepsilon}^2} \times \max_{\substack{0 \le i \le N_\varepsilon \\ m \in [\underline{m},\overline{m}]}} W(\sigma_m E_i) < \varepsilon. \tag{5.17}$$

- Step 3: construction in $[a_n, b_n]$. We start by splitting this interval into subintervals with integer "charge". This is done by induction by letting $t_0 = a_n$ and, t_k being given, letting t_{k+1} be the smallest $t \ge t_k + q_\varepsilon R_\varepsilon$ such that $\int_{t_{k-1}}^{t_k} m_0{}'(x)\, dx \in q_\varepsilon \mathbb{N}$. By (5.11) there exists $K \in \mathbb{N}$ such that $t_K = b_n$, and

$$K \le \frac{b_n - a_n}{q_\varepsilon R_\varepsilon} \le \frac{n(b-a)}{q_\varepsilon R_\varepsilon}. \tag{5.18}$$

Since $m_0{}' \ge \underline{m}$ in $[a_n, b_n]$, it is clear that $t_k - (q_\varepsilon R_\varepsilon + t_{k-1}) \le q_\varepsilon \underline{m}^{-1}$. To summarize and letting $I_k = [t_{k-1}, t_k]$, we thus have

$$|I_k| \in [q_\varepsilon R_\varepsilon, q_\varepsilon (R_\varepsilon + \underline{m}^{-1})], \qquad \int_{I_k} m_0{}'(x)\, dx \in q_\varepsilon \mathbb{N}. \tag{5.19}$$

In each I_k we will paste $n_{i,k}$ copies of a rescaled version of E_i, where

$$n_{i,k} = \left[\frac{q_\varepsilon (b_n - a_n)}{|I_k|} p_{i,k} \right] \qquad p_{i,k} = \tilde P \left(\frac{1}{n} I_k \times H^i_\varepsilon \right),$$

$[\cdot]$ denoting the integer part of a number. Because the first marginal of $\tilde P$ is the normalized Lebesgue measure on $\Sigma_{\underline{m}}$ and since $[a_n, b_n] \subset \Sigma_{\underline{m}} \subset [a,b]$, and $\cup_k I_k = [a_n, b_n]$, we have that

$$\frac{|I_k|}{n(b-a)} \le \sum_{i=1}^{N_\varepsilon} p_{i,k} \le \frac{|I_k|}{b_n - a_n},$$

and therefore $\sum_{i=1}^{N_\varepsilon} n_{i,k} \le q_\varepsilon$. Also, using in particular (5.17),

$$\sum_{i,k} \left| \frac{|I_k|}{q_\varepsilon(b_n - a_n)} n_{i,k} - p_{i,k} \right| \le \frac{N_\varepsilon}{q_\varepsilon} < \eta_0.$$

We divide I_k into q_ε subintervals with disjoint interiors, all having the same width $\in [R_\varepsilon, R_\varepsilon + \underline{m}^{-1}]$. Then for each $1 \le i \le N_\varepsilon$ we let $\mathcal{I}_{i,k}$ denote a family consisting of $n_{i,k}$ of these intervals. This doesn't necessarily exhaust I_k since $\sum_{i=1}^{N_\varepsilon} n_{i,k} \le q_\varepsilon$ so we let $n_{0,k} = q_\varepsilon - \sum_{i=1}^{N_\varepsilon} n_{i,k}$.

We define m_k to be the average of ${m_0}'$ over I_k. From (5.19) we have $m_k|I_k| \in q_\varepsilon\mathbb{N}$ hence for each $I \in \mathcal{I}_{i,k}$ we have $R := |m_k I| \in \mathbb{N}$, and $R \in [m_k R_\varepsilon, m_k(R_\varepsilon + \underline{m}^{-1})]$. We then apply Proposition 3.4 in I_R to the vector field E_i, which yields a "truncated" vector field $E_{i,I}$ defined in I_R, where $R = |m_k I|$. If $I \in \mathcal{I}_{0,k}$ we apply the same procedure with an arbitrary current $E_0 \in \mathcal{A}_1$ fixed with respect to all the parameters of the construction.

We then set

$$E_n^{(1)}(x) = \sigma_{1/m_k} E_{i,I}(x_I + \cdot)$$

on each interval $I \in \mathcal{I}_{i,k}$, where x_I is the center of I. The next step is to rectify the weight in $E_n^{(1)}$. For this we let $\mathcal{R}_k$ be the square $I_k \times (-|I_k|/2, |I_k|/2)$ and let H_k be the solution to

$$\begin{cases} -\Delta H_k = 2\pi({m_0}' - m_k) & \text{in} \mathcal{R}_k \\ \dfrac{\partial H_k}{\partial \nu} = 0 & \text{on } \partial \mathcal{R}_k. \end{cases}$$

From Lemma 4.4 applied with φ and m_0 equal to zero, and using the fact that m_0 is assumed to belong to $C^{0\frac{1}{2}}$, we have for any $q \in [1,4]$,

$$\int_{\mathcal{R}_k} |\nabla H_k|^q \le C_q |I_k|^2 \|{m_0}' - m_k\|^q_{L^\infty(I_k)} \le C_q |I_k|^2 \|m_0\|^q_{C^{0,\frac{1}{2}}} n^{-\frac{q}{2}}. \tag{5.20}$$

We then define

$$E_n^{(2)} = \begin{cases} \nabla H_k & \text{in } \mathcal{R}_k \\ 0 & \text{in } \bar{I}_k \backslash \mathcal{R}_k \end{cases}$$

$$E_n = E_n^{(1)} + E_n^{(2)} \quad \text{in } \bar{I}_k.$$

Using Lemma 4.6 and (5.20) we deduce using (5.19) that

$$W(E_n, \mathbf{1}_{\bar{I}_k}) \le W(E_n^{(1)}, \mathbf{1}_{\bar{I}_k}) + o_n(1), \quad \text{as } n \to \infty, \tag{5.21}$$

where $o_n(1)$ tends to zero as $n \to \infty$ and depends on $\varepsilon, \underline{m} > 0$ but not the interval I_k we are considering. Summing (5.20) for $1 \le k \le K$ and in view of (5.18) we find that for any $q \in [1,4]$

$$\int_{[a_n,b_n]\times\mathbb{R}} |E_n^{(1)} - E_n|^q \le C_{q,\varepsilon,\underline{m}} n^{1-\frac{q}{2}}. \tag{5.22}$$

On the other hand, in view of the construction and the result of Proposition 3.4 we have

$$W(E_n^{(1)}, \mathbf{1}_{\bar{I}_k}) \le |I_k| \left(\sum_{i=0}^{N_\varepsilon} \frac{n_{i,K}}{q_\varepsilon} W(\sigma_{m_K} J_i) + C\varepsilon \right). \tag{5.23}$$

Then, following the exact same arguments as in [SS2, Sec. 7] which we do not reproduce here (the only difference is that the rescaling factors $\sqrt{n}$ there should be replaced by n), thanks to (5.16)–(5.15)–(5.17) we find that we can choose C_1 in (5.17) such that

$$d_{\mathcal{B}}(\bar{P}, P') < C\varepsilon(|\log \varepsilon| + 1) \tag{5.24}$$

where

$$P' = \frac{1}{|\Sigma'_{\underline{m}}|} \sum_{k=1}^{K} \int_{I_k} \delta_{x_k} \otimes \delta_{\theta_\lambda E_n^{(1)}} \, d\lambda$$

and stands for $P^{(6)}$ in [SS2, Sec. 7]. Also, and again as in [SS2], since (5.22) holds, and from Lemma 5.2, we may replace $E_n^{(1)}$ with E_n at a negligible cost, more precisely for any large enough n we have

$$d_{\mathcal{B}}(\bar{P}, P'') < C\varepsilon(|\log \varepsilon| + 1) \tag{5.25}$$

where

$$P'' = \frac{1}{|\Sigma'_{\underline{m}}|} \sum_{k=1}^{K} \int_{I_k} \delta_{x_k} \otimes \delta_{\theta_\lambda E_n} \, d\lambda.$$

- *Step 3: construction in* $[b_n, nb]$. The construction in $[na, a_n]$ is exactly the same hence will be omitted. We claim that there exists E_n defined in $[b_n, nb] \times \mathbb{R}$ such that

$$\begin{cases} \operatorname{div} E_n = 2\pi(\sum_i \delta_{x_i} - {m_0}'\delta_{\mathbb{R}}) & \text{in } [b_n, nb] \times \mathbb{R} \\ E_n \cdot \vec{\nu} = 0 & \text{on } \partial([b_n, nb] \times \mathbb{R}) \end{cases} \tag{5.26}$$

and

$$W(E_n, \mathbf{1}_{[b_n, nb] \times \mathbb{R}}) \le Cn\left(\underline{m} + o_n(1)\right), \tag{5.27}$$

where C may depend on $\gamma, \overline{m}$ and ε. To prove this claim, let $s_0 = b_n$ and for every $l \ge 1$, let s_l be the smallest $s \ge s_{l-1}$ such that $\int_{s_{l-1}}^{s_l} {m_0}'(x)\, dx = 1$. Since (5.10) holds, this terminates at some $s_L = nb$ with $L = \int_{b_n}^{nb} {m_0}' \le \overline{m}|nb - b_n|$. We then set x_l to be the middle of $[s_{l-1}, s_l]$. We let u_l be the solution in the square $\mathcal{R}_l := [s_{l-1}, s_l] \times [-\frac{1}{2}(s_l - s_{l-1}), \frac{1}{2}(s_l - s_{l-1})]$

$$\begin{cases} -\Delta u_l = 2\pi(\delta_{x_l} - {m_0}'\delta_{\mathbb{R}}) & \text{in } \mathcal{R}_l \\ \dfrac{\partial u_l}{\partial \nu} = 0 & \text{on } \partial\mathcal{R}_l. \end{cases}$$

This equation is solvable since, by construction of the s_l's, the right-hand side has zero integral. Then for each l we let $E_n = -\nabla u_l$ in $\mathcal{R}_l$, and let $E_n = 0$ in $[b_n, nb] \times \mathbb{R} \setminus \cup_l \mathcal{R}_l$. Clearly E_n satisfies (5.26).

To estimate the energy of u_l we let $u_l = v_l + w_l$ where, letting $m_l = \fint_{[s_{l-1}, s_l]} {m_0}'$,

$$\begin{cases} -\Delta v_l = 2\pi(\delta_{x_i} - m_l\delta_{\mathbb{R}}) & \text{in } \mathcal{R}_l \\ \dfrac{\partial v_l}{\partial \nu} = 0 & \text{on } \partial\mathcal{R}_l, \end{cases}$$

$$\begin{cases} -\Delta w_l = 2\pi(m_l - {m_0}')\delta_{\mathbb{R}} & \text{in } \mathcal{R}_l \\ \dfrac{\partial w_l}{\partial \nu} = 0 & \text{on } \partial\mathcal{R}_l, \end{cases}$$

From Lemma 4.4 and Lemma 4.5 we find, choosing for instance $q = 4$ so that $q \in [1, 4]$ and $q' < 2$,

$$\int_{\mathcal{R}_l} |\nabla w_l|^q \le C(s_l - s_{l-1})^2 \left\|m_l - {m_0}'\right\|^q_{L^\infty([s_{l-1}, s_l])},$$

and

$$W(v_l, \mathbf{1}_{\mathcal{R}_l}) = C - \pi \log m_l, \qquad \int_{\mathcal{R}_l} |\nabla v_l|^{q'} \le C{m_l}^{q'-2}.$$

From (1.17) and Lemma 4.6, since $E_n = -(\nabla v_l + \nabla w_l)$ in $\mathcal{R}_l$, we have

$$\begin{aligned} (5.28) \quad W(E_n, \mathbf{1}_{\mathcal{R}_l}) \le C - \pi \log m_l + C \left\|m_l - {m_0}'\right\|_{L^\infty([s_{l-1}, s_l])} m_l^{1-\frac{2}{q'}} (s_l - s_{l-1})^{\frac{2}{q}} + \\ + C \left\|m_l - {m_0}'\right\|^2_{L^\infty([s_{l-1}, s_l])} (s_l - s_{l-1})^2. \end{aligned}$$

Using (1.18),

$$\left\|m_l - {m_0}'\right\|_{L^\infty([s_{l-1}, s_l])} \le C\|m_0\|_{C^{0,\frac{1}{2}}} \frac{(s_l - s_{l-1})^{\frac{1}{2}}}{\sqrt{n}}.$$

Replacing in (5.28) and letting $q = 4$ we deduce that

$$W(E_n, \mathbf{1}_{\mathcal{R}_l}) \le C - \pi \log m_l + C\left(\frac{s_l - s_{l-1}}{\sqrt{n}} + \frac{(s_l - s_{l-1})^3}{n}\right).$$

Then, summing with respect to l — using the fact that from (5.12) we have $\sum_l |s_{l+1} - s_l| \le Cn\overline{m}(1 + o_n(1))$, the fact that the integral over $[s_{l-1}, s_l]$ of ${m_0}'$ is 1 and that from (1.17) we have $(s_l - s_{l-1}) \le n^{\frac{1}{3}}$— we find

$$W(E_n, \mathbf{1}_{[b_n, nb]\times\mathbb{R}}) \le C\left(\int_{b_n}^{nb} {m_0}'(x) - {m_0}'(x)\log {m_0}'(x)\, dx + n o_n(1)\right) \le Cn(\underline{m} + o_n(1)),$$

since ${m_0}' - {m_0}' \log {m_0}'$ is bounded by a constant depending only on m_0 and using (5.12). This proves (5.27)

- *Step 4: conclusion.* Once the construction of E_n is completed, the proof of Proposition 3.1 essentially identical to that of [SS2, Proposition 4.1], which is its 2-dimensional equivalent, except that the scaling factor $\sqrt{n}$ there must be replaced by n. We only sketch the proof below and refer to the specific part of [SS2] for the details.

The test vector-field E_n has now been defined on all $[na, nb] \times \mathbb{R}$. It is extended by 0 outside, and is easily seen to satisfy the relation div $E_n = 2\pi(\nu'_n - {m_0}')$ for $\nu'_n = \sum_{i=1} \delta_{x'_i}$, a sum of Dirac masses on the real line. Combining (5.27) with (5.21), (5.23) and (5.6), we have

$$W(E_n, \mathbf{1}_{\mathbb{R}^2}) \le \sum_k |I_k| \left(\sum_{i=0}^{N_\varepsilon} \frac{n_{i,K}}{q_\varepsilon} W(\sigma_{m_K} E_i) + C\varepsilon + o_n(1)\right) + Cn(\underline{m} + o_n(1)).$$

Letting $n \to \infty$ and then $\underline{m} \to 0$, we see that the error term on the right-hand side can be made arbitrarily small, say smaller than $C\varepsilon$. On the other hand, the reasoning of [SS2], Step 2 in Paragraph 7.4, shows that

$$\sum_k |I_k| \left(\sum_{i=0}^{N_\varepsilon} \frac{n_{i,K}}{q_\varepsilon} W(\sigma_{m_K} E_i) \right) \le |\Sigma'| \int W(E)\, dP(x,E) + Cn\left(\varepsilon + o_n(1)\right),$$

so that taking n larger if necessary we obtain

$$\frac{1}{|\Sigma'|} W(E_n, \mathbf{1}_{\mathbb{R}^2}) \le \int W(E)\, dP(x,E) + C\varepsilon. \tag{5.29}$$

Then arguing as in Paragraph 7.4, Step 3 of [SS2] we obtain that, letting $(x_1, \cdots, x_n)$ be the rescalings to the original scale of the points x_i' i.e. $x_i = x_i'/n$, we have for n large enough

$$\limsup_{n\to\infty} \frac{1}{n} \left(w_n(x_1, \dots, x_n) - n^2 \mathcal{F}(\mu_0) + n \log n \right) \le \frac{|\Sigma|}{\pi} \int W(E)\, dP(x,E) + C\varepsilon.$$

Also letting P_n be the push-forward of $\frac{1}{|\Sigma|} dx\, |_\Sigma$ by the map $x \mapsto (x, E_n(nx + \cdot))$, it is easy to see that $d_{\mathcal{B}}(P'', P_n) < C\underline{m}$. In view of (5.13) and (5.25), and taking $\underline{m}$ small enough, for any given $\varepsilon > 0$, we can achieve

$$d_{\mathcal{B}}(P, P_n) < C\varepsilon.$$

This proves that items i) and ii) of Proposition 3.1 are satisfied by $(x_1, \dots, x_n)$ and E_n. Then, the perturbation argument of Paragraph 7.4, Step 4 in [SS2] shows that there exists $\delta > 0$ and for each n a subset $A_n \subset \mathbb{R}^n$ such that $|A_n| \ge n!(\delta/n)^n$ and such that for every $(y_1, \dots, y_n) \in A_n$ there exists a corresponding E_n satisfying (3.1) and (3.2). This concludes the proof of Proposition 3.1.

6 Proofs of the remaining theorems

6.1 Preliminary bounds on Z_n^β

The crucial fact is that there are good lower bounds for F_n. This follows from Proposition 4.1.

Even though we will not use the following result in the sequel, we state it to show how we can quickly derive a first upper bound on Z_n^β from what precedes.

Proposition 6.1. *We have*

$$\log K_n^\beta \le Cn\beta + n(\log|\Sigma| + o(1)) \tag{6.1}$$

and

$$\log Z_n^\beta \le -\beta n^2 \mathcal{F}(\mu_0) + \beta n \log n + Cn\beta + n(\log|\Sigma| + o(1)) \tag{6.2}$$

where $o(1) \to 0$ as $n \to \infty$ uniformly with respect to $\beta \ge \beta_0 > 0$, and C depends only on V.

The proof uses two lemmas.

Lemma 6.2. *For any* $\nu_n = \sum_{i=1}^n \delta_{x_i}$*, we have*

$$\widehat{F_n}(\nu_n) = \frac{1}{n\pi} \int_{\mathbb{R}^2} dg_{\nu_n} \tag{6.3}$$

where $\widehat{F_n}$ *is as in* (1.23) *and* g_{ν_n} *is the result of applying Proposition 4.1 to* ν_n*.*

Proof. This follows from (4.2) applied to χ_R, where χ_R is as in (1.7). If R is large enough then $\#\{p \in \mathrm{Supp}(\nu) \mid B(p, C) \cap \mathrm{Supp}(\nabla\bar{\chi}) \neq \varnothing\} = 0$ and therefore (4.2) reads

$$W(E_{\nu_n}, \chi_R) = \int \bar{\chi}_R \, dg_{\nu_n}.$$

Letting $R \to +\infty$ yields $W(E_{\nu_n}, \mathbf{1}_{\mathbb{R}^2}) = \int dg_{\nu_n}$ and the result, in view of (1.22). □

The following lemma has the same proof as in [SS2, Lemma 3.5].

Lemma 6.3. *Letting* ν_n *stand for* $\sum_{i=1}^n \delta_{x_i}$ *we have, for any constant* $\alpha > 0$ *and uniformly w.r.t.* $\beta \geq \beta_0 > 0$*,*

$$\lim_{n\to\infty} \left(\int_{\mathbb{R}^n} e^{-\alpha\beta n \int \zeta \, d\nu_n} \, dx_1 \dots dx_n \right)^{\frac{1}{n}} = |\Sigma|. \tag{6.4}$$

Proposition 6.1 is then proved exactly as in [SS2], Proposition 3.3. Note that a lower bound for Z_n^β was obtained in Corollary 3.3. To complete the proof of Theorem 3 it then suffices to prove the corresponding upper bound.

6.2 Lower bounds for $\widehat{F_n}$ and consequences

We start with a result that shows how $\widehat{F_n}$ controls the fluctuation $\nu_n - n\mu_0$.

Lemma 6.4. *Let* $\nu_n = \sum_{i=1}^n \delta_{x_i}$*. For any interval* I *of width* R *(possibly depending on* n*) and any* $1 < q < 2$*, we have*

$$\|\nu_n - n\mu_0\|_{W^{-1,q}(I)} \leq C_q (1 + R^2)^{\frac{1}{q} - \frac{1}{2}} n^{\frac{1}{2}} \left(\widehat{F_n}(\nu_n) + 1 \right)^{\frac{1}{2}}.$$

Here $W^{-1,q}$ *is the dual of the Sobolev space* $W_0^{1,q'}$ *with* $1/q + 1/q' = 1$*.*

Proof. In [SS2, Lemma 5.1], we have the following statement

$$\|\nu_n - n\mu_0 \delta_{\mathbb{R}}\|_{W^{-1,q}(B_R)} \leq C_q (1 + R^2)^{\frac{1}{q} - \frac{1}{2}} n^{\frac{1}{2}} \left(\widehat{F_n}(\nu_n) + 1 \right)^{\frac{1}{2}}.$$

The proof is based on [SeTi] which works in our one-dimensional context as well, thus the proof can be reproduced without change. It is immediate to deduce the result. □

We now turn to bounding from below $\widehat{F_n}$.

The proof is the same as in [SS2, Sec. 6], itself following the method of [SS1] based on the ergodic theorem. We just state the main ingredients.

Let $\{\nu_n\}_n$ and P_{ν_n} be as in the statement of Theorem 4. We need to prove that any subsequence of $\{P_{\nu_n}\}_n$ has a convergent subsequence and that the limit P is a $T_{\lambda(x)}$-invariant probability measure such that P-almost every (x, E) is such that $E \in \mathcal{A}_{m_0(x)}$ and (1.29) holds. Note that the fact that the first marginal of P is $dx_{|\Sigma}/|\Sigma|$ follows from the fact that, by definition, this is true of P_{ν_n}.

We thus take a subsequence of $\{P_{\nu_n}\}$ (which we don't relabel). We may assume that it has a subsequence, still denoted ν_n, which satisfies $\widehat{F_n}(\nu_n) \le C$, otherwise there is nothing to prove. This implies that ν_n is of the form $\sum_{i=1}^n \delta_{x_{i,n}}$. We let E_n denote the electric field and g_n the measures associated to ν_n as in Definition 4.3. As usual, ${\nu_n}' = \sum_{i=1}^n \delta_{nx_{i,n}}$.

A useful consequence of $\widehat{F_n}(\nu_n) \le C$ is that, using the result of Lemma 6.4, we have

$$\frac{1}{n}\nu_n \to \mu_0 \quad \text{on } \mathbb{R}. \tag{6.5}$$

We then set up the framework of Section 6.1 in [SS2]. We let $G = \Sigma$ and $X = \mathcal{M}_+ \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2) \times \mathcal{M}$, where $p \in (1, 2)$, where $\mathcal{M}_+$ denotes the set of positive Radon measures on $\mathbb{R}^2$ and $\mathcal{M}$ the set of those which are bounded below by the constant $-C_V := -C(\|m_0\|_\infty + 1)$ of Proposition 4.1, both equipped with the topology of weak convergence.

For $\lambda \in \mathbb{R}$ and abusing notation we let θ_λ denote both the translation $x \mapsto x + \lambda$ and the action

$$\theta_\lambda(\nu, E, g) = (\theta_\lambda \# \nu, E \circ \theta_\lambda, \theta_\lambda \# g).$$

Accordingly the action T^n on $\Sigma \times X$ is defined for $\lambda \in \mathbb{R}$ by

$$T^n_\lambda(x, \nu, E, g) = \left(x + \frac{\lambda}{n}, \theta_\lambda \# \nu, E \circ \theta_\lambda, \theta_\lambda \# g\right).$$

Then we let χ be a smooth nonnegative cut-off function with integral 1 and support in $[-1, 1]$ and define

$$\mathbf{f}_n(x, \nu, E, g) = \begin{cases} \dfrac{1}{\pi}\displaystyle\int_{\mathbb{R}^2} \chi(t)\, dg(t, s) & \text{if } (\nu, E, g) = \theta_{nx}({\nu_n}', E_n, g_n), \\ +\infty & \text{otherwise.} \end{cases} \tag{6.6}$$

Finally we let,

$$\mathbf{F}_n(\nu, E, g) = \fint_\Sigma \mathbf{f}_n\left(x, \theta_{xn}(\nu, E, g)\right)\, dx. \tag{6.7}$$

We have the following relation between $\mathbf{F}_n$ and $\widehat{F_n}$, as $n \to +\infty$ (see [SS2, Sec. 6]):

$$\mathbf{F}_n(\nu, E, g) \text{ is } \quad \begin{cases} \le \frac{1}{|\Sigma|}\widehat{F_n}(\nu_n) + o(1) & \text{if } (\nu, E, g) = ({\nu_n}', E_n, g_n) \\ = +\infty & \text{otherwise.} \end{cases} \tag{6.8}$$

The hypotheses in Section 6.1 of [SS2] are satisfied and applying the abstract result, Theorem 6 of [SS2], we conclude that letting Q_n denote the push-forward of the normalized Lebesgue measure on Σ by the map $x \mapsto (x, \theta_{nx}({\nu_n}', E_n, g_n))$, and $Q = \lim_n Q_n$, we have

$$\liminf_n \frac{1}{|\Sigma|}\widehat{F_n}(\nu_n) \ge \frac{1}{\pi}\int W(E)\, dQ(x, \nu, E, g) \tag{6.9}$$

and, Q-a.e. $(E, \nu) \in \mathcal{A}_{m_0(x)}$.

Now we let P_n (resp. P) be the marginal of Q_n (resp. Q) with respect to the variables (x, E). Then the first marginal of P is the normalized Lebesgue measure on E and P-a.e. we have $E \in \mathcal{A}_{m_0(x)}$, in particular

$$W(E) \geq \min_{\mathcal{A}_{m_0(x)}} W = m_0(x) \left(\min_{\mathcal{A}_1} W - \pi \log m_0(x) \right).$$

Integrating with respect to P and noting that since only x appears on the right-hand side we may replace P by its first marginal there, we find, in view of (1.27) that the lower bound (1.29) holds.

6.3 Proof of Theorems 4, 5 and 3, completed

As mentioned above, Part B of Theorem 4 is a direct consequence of Proposition 3.1, see Corollary 3.2.

Part C follows from the comparison of Parts A and B: for minimizers, the chains of inequalities (1.29) and (1.30) are in fact equalities and that $\frac{|\Sigma|}{\pi} \int W \, dP$ must be minimized hence equal to α, cf. (1.35). Since (1.29) follows from (6.9), the inequality (6.9) must also be an equality and moreover we must have $\lim_{n\to\infty} \int \zeta \, d\nu_n = 0$.

This completes the proof of Theorem 4. The proof of Theorems 5 and Theorem 3 is identical to [SS2] except again the replacement of the scaling $\sqrt{n}$ by n and $|A_n| \geq n!(\pi\delta^2/n)^n$ by $|A_n| \geq n!(\pi\delta/n)^n$.

6.4 Proof of Proposition 2.1

The result of Proposition 2.1 is a consequence of Proposition 3.4.

First, applying Part C of Theorem 4, there exists a translation-invariant measure P on $\Sigma \times L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that P-a.e. (x, E) is such that E minimizes W over $\mathcal{A}_{m_0(x)}$. Then, taking the push-forward of P under $(x, E) \mapsto \sigma_{1/m_0(x)} E$, we obtain a probability Q on $L^p_{\text{loc}}(\mathbb{R}^2, \mathbb{R}^2)$ such that Q-a.e. E minimizes W over $\mathcal{A}_1$.

Applying Lemma 5.4 to Q, we find that Q-a.e. E is such that $E \in \mathcal{A}_1$, such that (3.10) holds, and such that $W(E) = \min_{\mathcal{A}_1} W$. Choose on such E_0 and apply Proposition 3.4 to $G = \{E_0\}$. We find that for any given $\varepsilon > 0$ and any integer R large enough depending on ε, there exists E_R defined on $\bar{I}_R$ such that $E_R \cdot \vec{\nu} = 0$ on $\partial \bar{I}_R$ and $W(E_R, \mathbf{1}_{\bar{I}_R}) < R(W(E_0) + \varepsilon)$. This E_R can be extended periodically by letting $E_R(x + kR, y) = E_R(x, y)$ for any $k \in \mathbb{Z}$.

From the condition $E_R \cdot \vec{\nu} = 0$ on $\partial \bar{I}_R$ we find that, letting $\Lambda \subset I_R$ be the locations of the Dirac masses in div E_R, we have div $E_R = 2\pi \left(\sum_{p \in \Lambda_R} \delta_p - \delta_{\mathbb{R}} \right)$, where $\Lambda_R = \Lambda + R\mathbb{Z}$. Moreover

$$W(E_R) = \frac{W(E_R, \mathbf{1}_{\bar{I}_R})}{|I_R|} \leq W(E_0) + C\varepsilon.$$

There remains to make E_R curl-free. Following the proof of Corollary 4.4 of [SS1] we let $\tilde{E}_R = E_R + \nabla^\perp f_R$ in $\bar{I}_R$ where f_R solves $-\Delta f_R = \operatorname{curl} E_R$ in $\bar{I}_R$ and $f_R = 0$ on $\partial \bar{I}_R$. Then, div $\tilde{E}_R =$ div E_R and $\operatorname{curl} \tilde{E}_R = 0$ in $\bar{I}_R$. We can thus find H_R such that $\tilde{E}_R = \nabla H_R$ in $\bar{I}_R$. It also satisfies $\nabla H_R \cdot \vec{\nu} = \tilde{E}_R \cdot \vec{\nu} = E_R \cdot \vec{\nu} + \nabla^\perp f_R \cdot \vec{\nu} = 0$ on $\partial \bar{I}_R$. We may then extend H_R to a periodic function by even reflection, and take the final $\bar{E}_R$ to be ∇H_R. This procedure can only decrease the energy (arguing again as in [SS1, SS2]): we have $W(\tilde{E}_R, \mathbf{1}_{\bar{I}_R}) \leq W(E_R, \mathbf{1}_{\bar{I}_R})$

since

$$\int_{\bar I_R\backslash\cup B(p,\eta)}|\nabla H_R-\nabla^\perp f_R|^2-\int_{\bar I_R\backslash\cup B(p,\eta)}|\nabla H_R|^2$$
$$=-2\int_{\bar I_R\backslash\cup B(p,\eta)}\nabla H_R\cdot\nabla^\perp f_R+\int_{\bar I_R\backslash\cup B(p,\eta)}|\nabla f_R|^2.$$

It can be checked that the last two terms on the right-hand side converge as $\eta\to 0$ to the integrals over $\bar I_R$. Also integrating by parts, using the Jacobian structure and the boundary data, we have $\int_{\bar I_R}\nabla H_R\cdot\nabla^\perp f_R=0$. Therefore, letting $\eta\to 0$ in the above yields

$$W(E_R,\mathbf{1}_{\bar I_R})-W(\nabla H_R,\mathbf{1}_{\bar I_R})\geq 0.$$

We deduce that $W(\bar E_R)\leq W(E_R)\leq\min_{\mathcal{A}_1}W+C\varepsilon$, with $\bar E_R$ a $2R$-periodic (with respect to the variable x) test vector field belonging to $\mathcal{A}_1$. The result follows by a standard diagonal argument.

6.5 Consequences for deviations: proof of Theorem 6

The proof relies on the following proposition, whose proof is much shorter than in [SS2], due to the simpler nature of the one-dimensional geometry.

Proposition 6.5. *Let $\nu_n=\sum_{i=1}^n\delta_{x_i}$, and g_{ν_n} be as in Definition 4.3. For any $R>1$, for any $x_0\in\mathbb{R}$, denoting*

$$D(x_0,R)=\nu_n\left(B_{\frac{R}{n}}(x_0)\right)-n\mu_0\left(B_{\frac{R}{n}}(x_0)\right)$$

we have

$$\int_{B_{2R}(x_0')}dg_{\nu_n}\geq -CR+cD(x_0,R)^2\min\left(1,\frac{|D(x_0,R)|}{R}\right),$$

where $c>0$ and C depend only on V. [1]

Proof. Two cases can happen: either $D(x_0,R)\geq 0$ or $D(x_0,R)\leq 0$.

We start with the first case. Let us choose $\tau=\min\left(2,1+\frac{D(x_0,R)}{2R\|m_0\|_{L^\infty}}\right)$ and denote $T=\{r\in[R,\tau R],B_r(x_0')\cap\mathcal{B}_\rho=\varnothing\}$, where $\mathcal{B}_\rho$ is as in Proposition 4.1. By construction of $\mathcal{B}_\rho$ and since $\rho<\frac12$, we have $|T|\geq\frac12(\tau-1)R$. We then follow the method of "integrating over circles" introduced in [SS3]: let $\mathcal{C}$ denote $\{x\in B_{\tau R}(x_0')\backslash B_R(x_0'),|x-x_0'|\notin T\}$.

For any $r\in T$, since $\partial B_r(x_0')$ does not intersect $Supp(\nu_n')$, we have

$$\int_{\partial B_r(x_0')}E_{\nu_n}\cdot\nu=\int_{B_r(x_0')}\text{div } E_{\nu_n}=2\pi\nu_n'(B_r(x_0'))-\int_{B_r(x_0')}m_0\left(\frac{x}{n}\right)\delta_{\mathbb{R}}\tag{6.10}$$
$$\geq D(x_0,R)-2(\tau-1)R\|m_0\|_{L^\infty}\geq\frac12 D(x_0,R)$$

by assumption and by choice of τ. Moreover, for any $r\in T$, we have, by Cauchy-Schwarz,

$$\int_{\partial B_r(x_0')}|E_{\nu_n}|^2\geq\frac{1}{2\pi r}\left(\int_{\partial B_r(x_0')}E_{\nu_n}\cdot\vec\nu\right)^2\geq\frac{1}{8\pi r}D(x_0,R)^2.$$

---

[1] The condition $R>1$ could be replaced by $R>R_0$ for any $R_0>0$ at the expense of a constant c depending on R_0.

Integrating over T, using $|T| \geq \frac{1}{2}(\tau-1)R$, we have

$$\int_T \frac{dr}{r} \geq \int_{\tau R - \frac{1}{2}(\tau-1)R}^{\tau R} \frac{dr}{r} = -\log\left(1 - \frac{\tau-1}{2\tau}\right)$$

and thus

$$\int_{B_{\tau R}(x_0')\setminus \mathcal{B}_\rho} |E_{\nu_n}|^2 \geq cD(x_0,R)^2 \min\left(1, \frac{D(x_0,R)}{R\|m_0\|_{L^\infty}}\right),$$

for some $c>0$ depending only on $\|m_0\|_{L^\infty}$ hence on V. Inserting into (4.1), we are led to

$$\int_{B_{2R}(x_0')} g_{\nu_n} \geq -C(\|m_0\|_{L^\infty}+1)R + cD(x_0,R)^2 \min\left(1, \frac{D(x_0,R)}{R\|m_0\|_{L^\infty}}\right).$$

The case $D(x_0,R) \leq 0$ is essentially analogous.

□

We now proceed to the proof of Theorem 6, starting with (1.36). If $R > R_0$ and $|D(x_0', R)| \geq \eta R$ then from Proposition 6.5 and using the fact — from Proposition 4.1 — that g_{ν_n} is positive outside $\cup_{i=1}^n B(x_i', C)$ and that $g_{\nu_n} \geq -C$ everywhere, we deduce from (6.3) and (1.22), (1.23) that

$$F_n(\nu_n) \geq \frac{1}{n}\left(-CR + c\min(\eta^2,\eta^3)R^2\right) + 2\int \zeta\, d\nu_n. \tag{6.11}$$

Inserting into (3.4) we find

$$\mathbb{P}_n^\beta\left(\left|D(x_0',R)\right| \geq \eta R\right) \leq \frac{1}{K_n^\beta}\exp\left(C\beta R - c\beta\min(\eta^2,\eta^3)R^2\right)\int e^{-n\beta\int\zeta\, d\nu_n}\, dx_1\dots dx_n.$$

Then, using the lower bound (3.6) and Lemma 6.3 we deduce that if $\beta \geq \beta_0$ and n is large enough depending on β_0 then

$$\log \mathbb{P}_n^\beta\left(\left|D(x_0',R)\right| \geq \eta R\right) \leq -c\beta\min(\eta^2,\eta^3)R^2 + C\beta R + Cn\beta + Cn,$$

where $c, C > 0$ depend only on V. Thus (1.36) is established.

We next turn to (1.38). Arguing as above, from (6.3) we have $F_n(\nu_n) \geq -C + 2\int \zeta\, d\nu_n$. Splitting $2\int\zeta\, d\nu_n$ as $\int\zeta\, d\nu_n + \int\zeta\, d\nu_n$, inserting into (3.4) and using (3.6) we are led to

$$\mathbb{P}_n^\beta\left(\int \xi_n\, d\nu_n \geq \eta\right) \leq e^{-\frac{1}{2}n\beta\eta + Cn(\beta+1)}\int e^{-n\beta\int\zeta\, d\nu_n}\, dx_1\dots, dx_n,$$

where C depends only on V. Then, using Lemma 6.3 we deduce (1.38).

We finish with (1.37). Inserting the result of Lemma 6.4 into (3.4), we have, if I is an interval of width R/n

$$\mathbb{P}_n^\beta\left(\|\nu_n - n\mu_0\|_{W^{-1,q}(I)} \geq C_q\eta\sqrt{n}(1+R^2/n^2)^{\frac{1}{q}-\frac{1}{2}}\right) \leq \frac{1}{K_n^\beta}e^{-\frac{1}{2}n\beta\eta}\int e^{-n\beta\int\zeta\, d\nu}\, dx_1\dots dx_n.$$

Arguing as before and rearranging terms yields (1.37).

This concludes the proof of Theorem 6.

References

[ABF] G. Akemann, J. Baik, P. Di Francesco, *The Oxford Handbook of Random Matrix Theory*, Oxford University Press, 2011.

[AGZ] G. W. Anderson, A. Guionnet, O. Zeitouni, *An introduction to random matrices.* Cambridge University Press, 2010.

[AM] G. Alberti, S. Müller, A new approach to variational problems with multiple scales. *Comm. Pure Appl. Math.* **54**, no. 7 (2001), 761-825.

[AM] M. Aizenman, P. Martin, Structure of Gibbs States of one Dimensional Coulomb Systems, *Commun. Math. Phys.* **78** (1980), 99-116.

[Be] M. E. Becker, Multiparameter groups of measure-preserving transformations: a simple proof of Wiener's ergodic theorem. *Ann Probab.* **9**, No 3 (1981), 504–509.

[BG] G. Ben Arous, A. Guionnet, Large deviations for Wigner's law and Voiculescu's non-commutative entropy, *Probab. Theory Related Fields* **108** (1997), no. 4, 517–542.

[BSe] A. Borodin, S. Serfaty, Renormalized Energy Concentration in Random Matrices, to appear in *Comm. Math. Phys.*

[BEY1] P. Bourgade, L. Erdös, H.-T. Yau, Universality of general β-ensembles, `arXiv:0907.5605`.

[BEY2] P. Bourgade, L. Erdös, H. T. Yau, Bulk Universality of General β-ensembles with non-convex potential, `arXiv:1201.2283`.

[Br] A. Braides, *Γ-convergence for beginners*, Oxford University Press, 2002.

[BL] H. J. Brascamp, E. H. Lieb, In: *Functional integration and its applications,* Clarendon Press, 1975.

[CS] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian. *Comm. Partial Differential Equations* **32** (2007), no. 7-9, 1245–1260.

[D] P. Deift, *Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach.* Courant Lecture Notes in Mathematics, AMS, 1999.

[DG] P. Deift, D. Gioev, *Random matrix theory: invariant ensembles and universality.* Courant Lecture Notes in Mathematics, AMS, 2009.

[DE] I. Dumitriu, A. Edelman, Matrix models for beta ensembles, *J. Math. Phys.* **43** (2002), 5830–5847.

[Dy] F. Dyson, Statistical theory of the energy levels of a complex system. Part I, *J. Math. Phys.* **3**, 140–156 (1962); Part II, ibid. 157–165; Part III, ibid. 166–175

[EML] N. M. Ercolani, K. D. McLaughlin, Asymptotics of the partition function for random matrices via Riemann-Hilbert techniques and applications to graphical enumeration. *Int. Math. Res. Not.* **14** (2003), 755–820.

[EPRSY] L. Erdös, S. Péché, J. A. Ramirez, B. Schlein, H.T. Yau, Bulk universality for Wigner matrices, *Comm. Pure Appl. Math.* **63** (2010), 895–925.

[ESY] L. Erdös, B. Schlein, H.T. Yau, Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices. *Ann. Probab.* **37** (2009), no. 3, 815–852.

[For] P. J. Forrester, *Log-gases and random matrices.* London Mathematical Society Monographs Series, 34. Princeton University Press, 2010.

[Fro] O. Frostman, Potentiel d'équilibre et capacité des ensembles avec quelques applications à la théorie des fonctions. *Meddelanden Mat. Sem. Univ. Lund 3*, **115 s** (1935).

[GMS] A. Guionnet, E. Maurel-Segala, Combinatorial aspects of matrix models, *ALEA Lat. Am. J. Probab. Math. Stat.* **1** (2006), 241–279.

[Jo] K. Johansson, On fluctuations of eigenvalues of random Hermitian matrices, *Duke Math. J.* **91** (1998), no. 1, 151–204.

[Ko] W. König, Orthogonal polynomial ensembles in probability theory, *Probab. Surv.* **2** (2005), 385–447.

[Le1] A. Lenard, Exact statistical mechanics of a one-dimensional system with Coulomb forces. *J. Math. Phys.* **2** (1961), 682–693.

[Le2] A. Lenard, Exact statistical mechanics of a one-dimensional system with Coulomb forces. III. Statistics of the electric field, *J. Math. Phys.* **4** (1963), 533-543.

[Me] M. L. Mehta, *Random matrices. Third edition.* Elsevier/Academic Press, 2004.

[PAS] L. Pastur, S. Albeverio, M. Shcherbina, On the 1/n expansion for some unitary invariant ensembles of random matrices, *Comm. Math. Phys.* **224** (1), (2001), 271–305.

[SaTo] E. Saff, V. Totik, *Logarithmic potentials with external fields*, Springer-Verlag, 1997.

[SS1] E. Sandier, S. Serfaty, From the Ginzburg-Landau model to vortex lattice problems, *Comm. Math. Phys.* **313** (2012), no. 3, 635-743.

[SS2] E. Sandier, S. Serfaty, 2D Coulomb Gases and the Renormalized Energy, `arXiv:1201.3503.`

[SS3] E. Sandier, S. Serfaty, Global Minimizers for the Ginzburg-Landau Functional below the First critical Magnetic Field. *Annales Inst. H. Poincaré, Anal. non linéaire* **17** (2000), No 1, 119-145.

[SeTi] S. Serfaty, I. Tice, Lorentz space estimates for the Coulombian renormalized energy, *Commun. Contemp. Math.* **14** (2012), no. 4, 1250027, 23 pp.

[Si] B. Simon, The Christoffel-Darboux kernel, in "Perspectives in PDE, Harmonic Analysis and Applications," a volume in honor of V.G. Maz'ya's 70th birthday, *Proc. Symp. Pure Math.* **79** (2008), 295–335.

[TV] T. Tao, V. Vu, Random matrices: universality of local eigenvalue statistics up to the edge, *Comm. Math. Phys.* **298** (2) (2010) 549–572.

[VV] B. Valkó, B. Virág, Continuum limits of random matrices and the Brownian carousel. *Invent. Math.* **177** (2009), no. 3, 463–508.

[Wi] E. Wigner, Characteristic vectors of bordered matrices with infinite dimensions, *Ann. Math.* **62** (1955), 548–564.

Etienne Sandier
Université Paris-Est,
LAMA – CNRS UMR 8050,
61, Avenue du Général de Gaulle, 94010 Créteil. France
& Institut Universitaire de France
`sandier@u-pec.fr`

Sylvia Serfaty
UPMC Univ Paris 06, UMR 7598 Laboratoire Jacques-Louis Lions,
Paris, F-75005 France ;
CNRS, UMR 7598 LJLL, Paris, F-75005 France
& Courant Institute, New York University
251 Mercer st, NY NY 10012, USA
`serfaty@ann.jussieu.fr`